

Product datasheet for RC228162L3

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

COX11 (NM_001162861) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: COX11 (NM_001162861) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: COX11

Synonyms: COX11P

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC228162).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





st The last codon before the Stop codon of the ORF.

ACCN: NM_001162861

ORF Size: 690 bp





COX11 (NM_001162861) Human Tagged Lenti ORF Clone - RC228162L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001162861.1</u>, <u>NP 001156333.1</u>

RefSeq ORF: 693 bp Locus ID: 1353 Cytogenetics: 17q22

Protein Families: Transmembrane

Protein Pathways: Metabolic pathways, Oxidative phosphorylation

MW: 25.7 kDa

Gene Summary: Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain,

catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and

multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded


subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes a protein which is not a structural subunit, but may be a heme A biosynthetic enzyme involved in COX formation, according to the yeast mutant studies. However, the studies in Rhodobacter sphaeroides suggest that this gene is not required for heme A biosynthesis, but required for stable formation of the Cu(B) and magnesium centers of COX. This human protein is predicted to contain a transmembrane domain localized in the mitochondrial inner membrane. Multiple

pseudogene has been found on chromosome 6. [provided by RefSeq, Jun 2009]

transcript variants encoding different isoforms have been found for this gene. A related



Product images:



Circular map for RC228162L3