

Product datasheet for **RC228165**

COX11 (NM_001162862) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: COX11 (NM_001162862) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: COX11
Synonyms: COX11P
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >RC228165 representing NM_001162862
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGGAGGGCTCTGGCGTCTGGATGGAGGTGCGTTCCTTTCTGTGGCTGGCGCTGGATCCACCCTGGGT
CTCCAACCAGGGCTGCAGAGAGGGTAGAGCCGTTTCTTAGGCCAGAGTGGAGTGGGACAGGAGGTGCCGA
GAGAGGACTGAGGTGGCTTGGGACATGGAAGCGCTGCAGCCTTCGAGCCCGGCATCCAGCATTGCAGCCG
CCGCGCGGCCCTAAGAGCTCGAACCTTTCACACGCGCGCAGGAGGAGGAGCGCGCGGCAGAACAAGA
CGACCCTCACTTACGTGGCCGCTGTCGCGTGGGCATGCTGGGGCGTCTACGCTGCCGTACCCCTTTA
TCGGCTCTATTGCCAGACTACTGGACTTGGAGGATCAGCAGTTGCAGGTCATGCCTCAGACAAGATTGAA
AACATGGTGCCTGTTAAAGATCGAATCATTAAAATTAGCTTTAATGCAGATGTGCATGCAAGTCTCCAGT
GGAACCTTAGACCTCAGCAAACAGAAATATATGTGGTGCCAGGAGAGACTGCACTGGCGTTTTACAGAGC
TAAGAATCCTACTGACAAACCAGTAATTGGAATTTCTACATACAATATTGTTCCATTTGAAGCTGGACAG
TATTTCAATAAAATACAGGCTTCAAAGCTGCACAGAGTCTACGTTTTAGAGAGTTGGCACCTT

ACGCGTACGCGGGCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC228165 representing NM_001162862
Red=Cloning site Green=Tags(s)



MGGLWRPGWRCVPFCGWRWIHPGSPTRAAERVEPFLRPEWSGTGGAERGLRWLGTWKRCSLRARHPALQP
 PRRPKSSNPFFTRAQEEERRRQNKTTLYVAAVAVGMLGASYAAVPLYRLYCQTTGLGGSAVAGHASDKIE
 NMVPVKDRIIKISFNADVHASLQWNFRPQQTEIYVVPGETALAFYRAKNPTDKPVI GISTYNIIVPFEAGQ
 YFNKIQAASKLHRVYVLESWHL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



ACCN: NM_001162862

ORF Size: 693 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001162862.2](#)

RefSeq ORF: 696 bp

Locus ID: 1353

Cytogenetics: 17q22


Protein Families: Transmembrane

Protein Pathways: Metabolic pathways, Oxidative phosphorylation

MW: 26 kDa

Gene Summary: Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes a protein which is not a structural subunit, but may be a heme A biosynthetic enzyme involved in COX formation, according to the yeast mutant studies. However, the studies in *Rhodobacter sphaeroides* suggest that this gene is not required for heme A biosynthesis, but required for stable formation of the Cu(B) and magnesium centers of COX. This human protein is predicted to contain a transmembrane domain localized in the mitochondrial inner membrane. Multiple transcript variants encoding different isoforms have been found for this gene. A related pseudogene has been found on chromosome 6. [provided by RefSeq, Jun 2009]

Product images:



Circular map for RC228165