

Product datasheet for RC228219L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Triosephosphate isomerase (TPI1) (NM_001159287) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Triosephosphate isomerase (TPI1) (NM_001159287) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Triosephosphate isomerase

Synonyms: HEL-S-49; TIM; TPI; TPID

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001159287

ORF Size: 858 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC228219).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001159287.1, NP 001152759.1

 RefSeq ORF:
 861 bp

 Locus ID:
 7167

 UniProt ID:
 P60174

 Cytogenetics:
 12p13.31

Protein Pathways: Fructose and mannose metabolism, Glycolysis / Gluconeogenesis, Inositol phosphate

metabolism, Metabolic pathways





Triosephosphate isomerase (TPI1) (NM_001159287) Human Tagged ORF Clone Lentiviral Particle - RC228219L4V

MW: 30.6 kDa

Gene Summary: This gene encodes an enzyme, consisting of two identical proteins, which catalyzes the isomerization of glyceraldehydes 3-phosphate (G3P) and dihydroxy-acetone phosphate

(DHAP) in glycolysis and gluconeogenesis. Mutations in this gene are associated with

triosephosphate isomerase deficiency. Pseudogenes have been identified on chromosomes 1, 4, 6 and 7. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Apr

2009]