

## Product datasheet for RC228362L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RRM2 (NM\_001165931) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** RRM2 (NM\_001165931) Human Tagged ORF Clone Lentiviral Particle

Symbol: RRM2

Synonyms: C2orf48; R2; RR2; RR2M

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_001165931

ORF Size: 1347 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC228362).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** NM 001165931.1, NP 001159403.1

 RefSeq ORF:
 1350 bp

 Locus ID:
 6241

 UniProt ID:
 P31350

Cytogenetics: 2p25.1

**Protein Families:** Druggable Genome

**Protein Pathways:** Glutathione metabolism, Metabolic pathways, p53 signaling pathway, Purine metabolism,

Pyrimidine metabolism





## RRM2 (NM\_001165931) Human Tagged ORF Clone Lentiviral Particle - RC228362L2V

MW:

50.9 kDa

**Gene Summary:** 

This gene encodes one of two non-identical subunits for ribonucleotide reductase. This reductase catalyzes the formation of deoxyribonucleotides from ribonucleotides. Synthesis of the encoded protein (M2) is regulated in a cell-cycle dependent fashion. Transcription from this gene can initiate from alternative promoters, which results in two isoforms that differ in the lengths of their N-termini. Related pseudogenes have been identified on chromosomes 1 and X. [provided by RefSeq, Sep 2009]