

Product datasheet for RC228565L4

UROC1 (NM_001165974) Human Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	UROC1 (NM_001165974) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	UROC1
Synonyms:	HMFN0320; UROCD
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC228565).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	


Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.



[View online »](#)

Plasmid Map:


ACCN: NM_001165974

ORF Size: 2208 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001165974.1](#)

RefSeq ORF: 2211 bp

Locus ID: 131669

UniProt ID: [Q96N76](#)

Cytogenetics: 3q21.3

Protein Pathways: Histidine metabolism, Metabolic pathways

MW: 80.7 kDa

Gene Summary: This gene encodes an enzyme involved in the second step of histidine catabolism, metabolizing urocanic acid to formiminoglutamic acid. Deficiency of this enzyme results in urocanic aciduria, and is an apparent cause of mental retardation. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Apr 2021]