

## Product datasheet for **RC228595L3V**

### **ALOXE3 (NM\_001165960) Human Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	ALOXE3 (NM_001165960) Human Tagged ORF Clone Lentiviral Particle
Symbol:	ALOXE3
Synonyms:	ARCI3; E-LOX; eLOX-3; eLOX3
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001165960
ORF Size:	2529 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC228595).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_001165960.1</a> , <a href="#">NP_001159432.1</a>
RefSeq ORF:	2532 bp
Locus ID:	59344
UniProt ID:	<a href="#">Q9BYJ1</a>
Cytogenetics:	17p13.1
Protein Families:	Druggable Genome
MW:	95 kDa



[View online »](#)

**Gene Summary:**

This gene is a member of the lipoxygenase family, which are catabolized by arachidonic acid-derived compounds. The encoded enzyme is a hydroperoxide isomerase that synthesizes a unique type of epoxy alcohol (8R-hydroxy-11R,12R-epoxyeicosa-5Z,9E,14Z-trienoic acid) from 12R-hydroperoxyeicosatetraenoic acid (12R-HPETE). This epoxy alcohol can activate the nuclear receptor peroxisome proliferator-activated receptor alpha (PPARalpha), which is implicated in epidermal differentiation. Loss of function of the enzyme encoded by this gene results in ichthyosis, implicating the function of this gene in the differentiation of human skin. This gene is part of a cluster of lipoxygenase genes on 17p13.1. Mutations in this gene result in nonbullous congenital ichthyosiform erythroderma (NCIE). Multiple transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Sep 2009]