

### Product datasheet for RC228734L3V

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Tryptophan rich protein (WRB) (NM\_001146218) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Tryptophan rich protein (WRB) (NM\_001146218) Human Tagged ORF Clone Lentiviral Particle

Symbol: Tryptophan rich protein

**Synonyms:** CHD5; WRB

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001146218

ORF Size: 522 bp

**ORF Nucleotide** 

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC228734).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001146218.1</u>

 RefSeq ORF:
 423 bp

 Locus ID:
 7485

 UniProt ID:
 000258

Cytogenetics: 21q22.2

**Protein Families:** Transmembrane

**MW:** 19.78 kDa





# Tryptophan rich protein (WRB) (NM\_001146218) Human Tagged ORF Clone Lentiviral Particle – RC228734L3V

#### **Gene Summary:**

This gene is located in the candidate region for congenital heart disease (CHD) in Down syndrome (DS). It encodes a basic protein that functions as a receptor that promotes insertion of tail-anchored proteins in the endoplasmic reticulum membrane. This gene is located at a maternally-methylated differentially methylated region (DMR); however, its transcription may be biallelic, not imprinted. Alternative splicing results in different transcript variants. A pseudogene has been defined on chromosome 4. [provided by RefSeq, Apr 2017]