

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC228898L3V

ADA2a (TADA2A) (NM_001166105) Human Tagged ORF Clone Lentiviral Particle

Product data:

| Product Type: | Lentiviral Particles |
|------------------------------|---|
| Product Name: | ADA2a (TADA2A) (NM_001166105) Human Tagged ORF Clone Lentiviral Particle |
| Symbol: | ADA2a |
| Synonyms: | ADA2; ADA2A; hADA2; KL04P; TADA2L |
| Mammalian Cell Selection: | Puromycin |
| Vector: | pLenti-C-Myc-DDK-P2A-Puro (PS100092) |
| Tag: | Myc-DDK |
| ACCN: | NM_001166105 |
| ORF Size: | 1329 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(RC228898). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u> |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | <u>NM 001166105.1</u> |
| RefSeq Size: | 1678 bp |
| RefSeq ORF: | 1332 bp |
| Locus ID: | 6871 |
| UniProt ID: | <u>075478</u> |
| Cytogenetics: | 17q12 |
| Protein Families: | Transcription Factors |
| MW: | 51.5 kDa |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CRIGENE ADA2a (TADA2A) (NM_001166105) Human Tagged ORF Clone Lentiviral Particle – RC228898L3V

Gene Summary:Many DNA-binding transcriptional activator proteins enhance the initiation rate of RNA
polymerase II-mediated gene transcription by interacting functionally with the general
transcription machinery bound at the basal promoter. Adaptor proteins are usually required
for this activation, possibly to acetylate and destabilize nucleosomes, thereby relieving
chromatin constraints at the promoter. The protein encoded by this gene is a transcriptional
activator adaptor and has been found to be part of the PCAF histone acetylase complex.
Several alternatively spliced transcript variants encoding different isoforms of this gene have
been described, but the full-length nature of some of these variants has not been
determined. [provided by RefSeq, Oct 2009]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US