

Product datasheet for RC228962L4

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FATP2 (SLC27A2) (NM_001159629) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: FATP2 (SLC27A2) (NM_001159629) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: FATP2

Synonyms: ACSVL1; FACVL1; FATP2; hFACVL1; HsT17226; VLACS; VLCS

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC228962).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:






^{*} The last codon before the Stop codon of the ORF.





Plasmid Map:



ACCN: NM_001159629

ORF Size: 1701 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001159629.1</u>, <u>NP 001153101.1</u>

RefSeq ORF: 1704 bp **Locus ID:** 11001



FATP2 (SLC27A2) (NM_001159629) Human Tagged Lenti ORF Clone - RC228962L4

 UniProt ID:
 O14975

 Cytogenetics:
 15q21.2

Protein Families: Transmembrane

Protein Pathways: PPAR signaling pathway

MW: 64.4 kDa

Gene Summary: The protein encoded by this gene is an isozyme of long-chain fatty-acid-coenzyme A ligase

family. Although differing in substrate specificity, subcellular localization, and tissue distribution, all isozymes of this family convert free long-chain fatty acids into fatty acyl-CoA esters, and thereby play a key role in lipid biosynthesis and fatty acid degradation. This isozyme activates long-chain, branched-chain and very-long-chain fatty acids containing 22 or more carbons to their CoA derivatives. It is expressed primarily in liver and kidney, and is present in both endoplasmic reticulum and peroxisomes, but not in mitochondria. Its decreased peroxisomal enzyme activity is in part responsible for the biochemical pathology in

X-linked adrenoleukodystrophy. Alternatively spliced transcript variants encoding different

isoforms have been found for this gene. [provided by RefSeq, Apr 2009]