

Product datasheet for RC229049L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GLI1 (NM_001160045) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: GLI1 (NM_001160045) Human Tagged ORF Clone Lentiviral Particle

Symbol: GLI1

Synonyms: GLI; PAPA8; PPD1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001160045

ORF Size: 3321 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC229049).

OTI Disclaimer:

Cytogenetics:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001160045.1</u>, <u>NP 001153517.1</u>

12q13.3

 RefSeq Size:
 3414 bp

 RefSeq ORF:
 2937 bp

 Locus ID:
 2735

 UniProt ID:
 P08151

Protein Families: Adult stem cells, Cancer stem cells, Druggable Genome, Embryonic stem cells, ES Cell

Differentiation/IPS, Stem cell relevant signaling - DSL/Notch pathway, Transcription Factors





GLI1 (NM_001160045) Human Tagged ORF Clone Lentiviral Particle - RC229049L3V

Protein Pathways: Basal cell carcinoma, Hedgehog signaling pathway, Pathways in cancer

MW: 117.9 kDa

Gene Summary: This gene encodes a member of the Kruppel family of zinc finger proteins. The encoded

transcription factor is activated by the sonic hedgehog signal transduction cascade and regulates stem cell proliferation. The activity and nuclear localization of this protein is negatively regulated by p53 in an inhibitory loop. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2009]