

Product datasheet for RC229531L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Neurokinin B (TAC3) (NM 001178054) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Neurokinin B (TAC3) (NM_001178054) Human Tagged ORF Clone Lentiviral Particle

Symbol: TAC3

Synonyms: HH10; LncZBTB39; NK3; NKB; NKNB; PRO1155; ZNEUROK1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001178054

ORF Size: 309 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC229531).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001178054.1

RefSeq ORF: 312 bp Locus ID: 6866

UniProt ID: Q9UHF0

Cytogenetics: 12q13.3

Protein Families: Druggable Genome, Secreted Protein

MW: 11.8 kDa





Neurokinin B (TAC3) (NM_001178054) Human Tagged ORF Clone Lentiviral Particle – RC229531L4V

Gene Summary:

This gene encodes a member of the tachykinin family of secreted neuropeptides. The encoded preproprotein is proteolytically processed to generate the mature peptide, which is primarily expressed in the central and peripheral nervous systems and functions as a neurotransmitter. This peptide is the ligand for the neurokinin-3 receptor. This protein is also expressed in the outer syncytiotrophoblast of the placenta and may be associated with pregnancy-induced hypertension and pre-eclampsia. Mutations in this gene are associated with normosmic hypogonadotropic hypogonadism. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Feb 2016]