

## Product datasheet for RC229608

### HDAC8 (NM\_001166422) Human Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** HDAC8 (NM\_001166422) Human Tagged ORF Clone  
**Tag:** Myc-DDK  
**Symbol:** HDAC8  
**Synonyms:** CDA07; CDLS5; HD8; HDACL1; KDAC8; MRXS6; RPD3; WTS  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Cell Selection:** Neomycin  
**ORF Nucleotide Sequence:** >RC229608 representing NM\_001166422  
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCCGCGATCGCC

ATGGAGGAGCCGGAGGAACCGCGGACAGTGGCAGTCGCTGGTCCCGTTTATATCTATAGTCCCAGT  
ATGTCAGTATGTGTGACTCCCTGGCCAAGATCCCCAACGGCCAGTATGGTGCATTCTTTGATTGAAGC  
ATATGCACTGCATAAGCAGATGAGGATAGTTAAGCCTAAAGTGGCCTCCATGGAGGAGATGGCCACCTTC  
CACACTGATGCTTATCTGCAGCATCTCCAGAAGGTCAGCCAAGAGGGCGATGATGATCATCCGGACTCCA  
TAGAATATGGGCTAGGTTATGACTGCCAGCCACTGAAGGGATATTTGACTATGCAGCAGCTATAGGAGG  
GGCTACGATCACAGCTGCCCAATGCCTGATTGACGGAATGTGCAAAGTAGCAATTAAGTGGTCTGGAGGG  
TGGCATCATGCAAAGAAAGAGACGTGTGTATGTGGCACTTTACAAGGCATTC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >RC229608 representing NM\_001166422  
Red=Cloning site Green=Tags(s)

MEEPEPADSQSLVPVYIYSPEYVSMCDLAKIPKRASMVHSLIEAYALHKQMRIVKPKVASMEEMATF  
HTDAYLQHLQKVSQEGDDHPDSIEYGLGYDCPATEGIFDYAAAIGGATITAAQCLIDGMCKVAINWSSG  
WHHAKKETCVYVALYKAF

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** SgfI-MluI




[View online »](#)

## Cloning Scheme:



## Plasmid Map:



ACCN: NM\_001166422

ORF Size: 474 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u><a href="#">NM_001166422.2</a></u>
<b>RefSeq ORF:</b>	477 bp
<b>Locus ID:</b>	55869
<b>UniProt ID:</b>	<u><a href="#">Q9BY41</a></u>
<b>Cytogenetics:</b>	Xq13.1
<b>Protein Families:</b>	Druggable Genome, Transcription Factors
<b>MW:</b>	17.9 kDa
<b>Gene Summary:</b>	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class I of the histone deacetylase family. It catalyzes the deacetylation of lysine residues in the histone N-terminal tails and represses transcription in large multiprotein complexes with transcriptional co-repressors. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2009]