

## **Product datasheet for RC229677**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Galectin 3 (LGALS3) (NM\_001177388) Human Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Galectin 3 (LGALS3) (NM\_001177388) Human Tagged ORF Clone

Tag: Myc-DDK
Symbol: Galectin 3

Synonyms: CBP35; GAL3; GALBP; GALIG; L31; LGALS2; MAC2

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >RC229677 representing NM\_001177388
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

 ${\tt TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC}$ 

GCCGCGATCGCC

TTTTTACATTCATCAATATCCCTCTTGTAAGTCATCTACT

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >RC229677 representing NM\_001177388

Red=Cloning site Green=Tags(s)

MADNFSLHDALSGSGNPNPQGWPGAWGNQPAGAGGYPGASYPGAYPGQAPPGAYPGQAPPGAYPGAYAGAYPGAPAGYPGAPAGYPSGPGAYPSSGQPSATGAYPATGPYGAPAGPLIVPYNLPLPGGVVPRMLITILGTVKP

NANRIALDFQRGNDVAFHFNPRFNENNRRVIVCTYMCKGFMFTVSENFYIHQYPSCKSST

TRTRPLEQKLISEEDLAANDILDYKDDDDKV







Chromatograms: <a href="https://cdn.origene.com/chromatograms/ja3211\_e01.zip">https://cdn.origene.com/chromatograms/ja3211\_e01.zip</a>

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_001177388

ORF Size: 600 bp

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customer.com">customer.com</a> or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 001177388.1</u>, <u>NP 001170859.1</u>

RefSeq Size: 744 bp
RefSeq ORF: 603 bp
Locus ID: 3958
Cytogenetics: 14q22.3

**Protein Families:** Secreted Protein

**MW:** 20.5 kDa


**Gene Summary:** This gene encodes a member of the galectin family of carbohydrate binding proteins.

Members of this protein family have an affinity for beta-galactosides. The encoded protein is characterized by an N-terminal proline-rich tandem repeat domain and a single C-terminal carbohydrate recognition domain. This protein can self-associate through the N-terminal domain allowing it to bind to multivalent saccharide ligands. This protein localizes to the extracellular matrix, the cytoplasm and the nucleus. This protein plays a role in numerous cellular functions including apoptosis, innate immunity, cell adhesion and T-cell regulation. The protein exhibits antimicrobial activity against bacteria and fungi. Alternate splicing results

in multiple transcript variants.[provided by RefSeq, Oct 2014]



## **Product images:**



Circular map for RC229677