

## Product datasheet for RC229682L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MYD88 (NM\_001172569) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** MYD88 (NM\_001172569) Human Tagged ORF Clone Lentiviral Particle

Symbol: MYD88

Synonyms: IMD68; MYD88D

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001172569

ORF Size: 612 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC229682).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** NM 001172569.1, NP 001166040.1

 RefSeq ORF:
 576 bp

 Locus ID:
 4615

 UniProt ID:
 Q99836

Cytogenetics: 3p22.2

**Protein Families:** Druggable Genome

**Protein Pathways:** Apoptosis, Toll-like receptor signaling pathway

MW: 22.6 kDa







## **Gene Summary:**

This gene encodes a cytosolic adapter protein that plays a central role in the innate and adaptive immune response. This protein functions as an essential signal transducer in the interleukin-1 and Toll-like receptor signaling pathways. These pathways regulate that activation of numerous proinflammatory genes. The encoded protein consists of an N-terminal death domain and a C-terminal Toll-interleukin1 receptor domain. Patients with defects in this gene have an increased susceptibility to pyogenic bacterial infections. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Feb 2010]