

Product datasheet for **RC229743**

Tissue Factor (F3) (NM_001178096) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Tissue Factor (F3) (NM_001178096) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Tissue Factor
Synonyms:	CD142; TF; TFA
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	<p>>RC229743 representing NM_001178096</p> <p>Red=Cloning site Blue=ORF Green=Tags(s)</p>

TTTGTGAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGAGACCCCTGCCTGGCCCCGGGTCCCGCGCCCCGAGACCGCCGTCGCTCGGACGCTCCTGCTCGGCT
 GGGTCTTCGCCCAGGTGGCCGGCGCTTCAGGCACTACAAATACTGTGGCAGCATATAATTTAACTTGAA
 ATCAACTAATTTCAAGACAATTTGGAGTGGGAACCCAAACCGTCAATCAAGTCTACACTGTTCAAATA
 AGCACTAAGTCAGGAGATTGAAAAGCAAATGCTTTACACAACAGACACAGAGTGTGACCTCACCGACG
 AGATTGTGAAGGATGTGAAGCAGACGTAATGGCACGGGTCTTCTCTACCCGGCAGGGAATGTGGAGAG
 CACCGGTTCTGCTGGGGAGCCTCTGTATGAGAATCCCCAGAGTTCACACCTTACCTGGAGACAAACCTC
 GGACAGCCAACAATTCAGAGTTTGAACAGGTGGGAACAAAAGTGAATGTGACCGTAGAAGATGAACGGA
 CTTTAGTCAGAAGGAACAACACTTTCTAAGCCTCCGGGATGTTTTGGCAAGGACTTAATTTATACACT
 TTATTATTGGAATCTTCAAGTTCAGGAAAGAAATATTCTACATCATTGGAGCTGTGGTATTGTGGTCA
 TCATCCTTGTCATCATCTGGCTATATCTCTACACAAGTGTAGAAAGGCAGGAGTGGGGCAGAGCTGGAA
 GGAGAACTCCCCAC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA


[View online »](#)

Protein Sequence: >RC229743 representing NM_001178096
 Red=Cloning site Green=Tags(s)



METPAWPRVPRPETAVARTLLLGWVFAQVAGASGTTNTVAAYNLTKWSTNFKTILEWEPKPVNQVYTVQI
 STKSGDWKSKCFYTTDTECDLTDEIVKDKVQTYLARVFSYPAGNVESTGSAGEPLYENSPEFTPYLETNL
 GQPTIQSFEQVGTKVNVTVEDERTLVRRNNTFLSLRDVFGKDLIYTLYYWKSSSGKKYKSTLELWYLWS
 SSLSSSWLYLYTYSVEROEWGRAGRTPH

TRTRPLEOKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM 001178096

ORF Size: 714 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM_001178096.1, NP_001171567.1</u>
RefSeq ORF:	717 bp
Locus ID:	2152
UniProt ID:	<u>P13726</u>
Cytogenetics:	1p21.3
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Complement and coagulation cascades
MW:	27.6 kDa
Gene Summary:	<p>This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces, for example, on monocytes. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. Platelets and monocytes have been shown to express this coagulation factor under procoagulatory and proinflammatory stimuli, and a major role in HIV-associated coagulopathy has been described. Platelet-dependent monocyte expression of coagulation factor III has been described to be associated with Coronavirus Disease 2019 (COVID-19) severity and mortality. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Aug 2020]</p>

Product images:



Circular map for RC229743