

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC229831L3V

## HDAC8 (NM\_001166418) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	HDAC8 (NM_001166418) Human Tagged ORF Clone Lentiviral Particle
Symbol:	HDAC8
Synonyms:	CDA07; CDLS5; HD8; HDACL1; KDAC8; MRXS6; RPD3; WTS
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001166418
ORF Size:	858 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC229831).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001166418.1</u>
RefSeq ORF:	861 bp
Locus ID:	55869
UniProt ID:	<u>Q9BY41</u>
Cytogenetics:	Xq13.1
Protein Families:	Druggable Genome, Transcription Factors
MW:	32.4 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Histones play a critical role in transcriptional regulation, cell cycle progression, and<br/>developmental events. Histone acetylation/deacetylation alters chromosome structure and<br/>affects transcription factor access to DNA. The protein encoded by this gene belongs to class I<br/>of the histone deacetylase family. It catalyzes the deacetylation of lysine residues in the<br/>histone N-terminal tails and represses transcription in large multiprotein complexes with<br/>transcriptional co-repressors. Multiple transcript variants encoding different isoforms have<br/>been found for this gene. [provided by RefSeq, Oct 2009]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US