

Product datasheet for RC229874

VEGFA (NM_001171622) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
 Product Name: VEGFA (NM_001171622) Human Tagged ORF Clone
 Tag: Myc-DDK
 Symbol: VEGFA
 Synonyms: MVCD1; VEGF; VPF
 Mammalian Cell Selection: Neomycin
 Vector: pCMV6-Entry (PS100001)
 E. coli Selection: Kanamycin (25 ug/mL)
 ORF Nucleotide Sequence: >RC229874 representing NM_001171622
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

CTGACGGACAGACAGACAGACACCGCCCCAGCCCCAGCTACCACCTCCTCCCCGGCCGGCGGGACAG
 TGGACGCGGGCGGAGCCGCGGGCAGGGGCCGAGCCCGCCGAGGGCGGGGTGGAGGGGTCCGGGC
 TCGCGCGTCCGACTGAACTTTTCGTCCAACCTCTGGGCTGTTCTCGCTTCGGAGGAGCCGTGGTCCG
 GCGGGGAAGCCGAGCCGAGCGGAGCCGCGAGAAGTGCTAGCTCGGGCCGGGAGGAGCCGACGCCGAGG
 AGGGGAGGAGGAAGAAGAGAAGGAAGAGAGAGGGGCCGAGTGGCGACTCGGCGCTCGAAGCCGGG
 CTATGGACGGGTGAGGCGCGGTGTGCGCAGACAGTGTCCAGCCGCGCGCTCCCCAGGCCCTGGCC
 CGGGCCTCGGGCCGGGAGGAAGTAGCTCGCCGAGGCGCCGAGGAGAGCGGGCCGCCACAGCCGA
 GCCGGAGAGGGAGCGGAGCCGCGCCGCCCCGGTCCGGCCTCCGAAACCATGAACCTTCTGCTGTCTTG
 GGTGCAATTGGAGCCTTGCTTGCTCTACCTCCACCATGCCAAGTGGTCCCAGGCTGCACCCATGGCA
 GAAGGAGGAGGGCAGAATCATCACGAAGTGGTGAAGTTCATGGATGTCTATCAGCGCAGCTACTGCCATC
 CAATCGAGACCCTGGTGGACATCTCCAGGAGTACCCTGATGAGATCGAGTACATCTTCAAGCCATCCTG
 TGTGCCCTGATGCGATGCGGGGCTGCTGCAATGACGAGGGCCGGAGTGTGTGCCACTGAGGAGTCC
 AACATCACCATGCAGATTATGCGGATCAAACCTACCAAGGCCAGCAGATAGGAGAGATGAGCTTCTAC
 AGCACAACAAATGTGAATGCAGATGTGACAAGCCGAGGCGG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC229874 representing NM_001171622
Red=Cloning site Green=Tags(s)

LTDRQTD TAPSPSYHLLPGRRTVDAAASRGQGPEPAPGGGVEGVGARGVALKLFVQLLGCSRFGGAVVR
 AGEAEP SGAARSASSGREEPQPEEGEEEEKEEERGPQWRLGARKPGSWTGEAAVCADSAPAARAPQALA
 RASGRGGRVARRGAEESGPPHSPRRGSASRAGPGRASETMNFLLSWVHWSLALLLLYHHAKWSQAAPMA
 EGGGQNHHEVVKFMDVYQRSYCHPIETLVDFIQEYYPDEIEYIFKPSCVPLMRCGGCCNDEGLECVPTES
 NITMQIMRIKPHQGHIGMSFLQHMKCECRCDKPRR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk8051_d11.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_001171622

ORF Size: 951 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.


Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001171622.2
RefSeq ORF:	954 bp
Locus ID:	7422
UniProt ID:	P15692
Cytogenetics:	6p21.1
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	Bladder cancer, Cytokine-cytokine receptor interaction, Focal adhesion, mTOR signaling pathway, Pancreatic cancer, Pathways in cancer, Renal cell carcinoma, VEGF signaling pathway
MW:	34.8 kDa
Gene Summary:	<p>This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome, also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site. The levels of VEGF are increased during infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), thus promoting inflammation by facilitating recruitment of inflammatory cells, and by increasing the level of angiopoietin II (Ang II), one of two products of the SARS-CoV-2 binding target, angiotensin-converting enzyme 2 (ACE2). In turn, Ang II facilitates the elevation of VEGF, thus forming a vicious cycle in the release of inflammatory cytokines. [provided by RefSeq, Jun 2020]</p>

Product images:



Circular map for RC229874



HEK293T cells were transfected with the overexpression plasmids of 9 VEGF isoforms (from left to right: VEGF206, Cat# [RC223789]; VEGF189, Cat# [RC229706]; L-VEGF165, Cat# [RC223884]; VEGF165, Cat# [RC229662]; L-VEGF121, Cat# [RC222129]; VEGF iso18, Cat# RC229874; VEGF183, Cat# [RC229686]; VEGF145, Cat# [RC231952]; L-VEGF189, Cat# [RC224244]) for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-flag antibody (Cat# [TA180144], 1:1000) or anti-VEGFA mouse monoclonal antibody. (Cat# [TA500289], 1:500)