

Product datasheet for RC229936

CXCR2 (NM_001168298) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	CXCR2 (NM_001168298) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	CXCR2
Synonyms:	CD182; CDw128b; CMKAR2; IL8R2; IL8RA; IL8RB
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC229936 representing NM_001168298 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGAAGATTTTAAACATGGAGAGTGACAGCTTTGAAGATTTCTGGAAAGGTGAAGATCTTAGTAATTACA
GTTACAGCTCTACCCTGCCCCCTTTCTACTAGATGCCGCCCATGTGAACCAGAATCCCTGGAAATCAA
CAAGTATTTTGTGGTCATTATCTATGCCCTGGTATTCCTGCTGAGCCTGCTGGGAACTCCCTCGTGATG
CTGGTCATCTTATACAGCAGGGTCGGCCGCTCCGTCACCTGATGTCTACCTGCTGAACCTAGCCTTGCCG
ACCTACTCTTTGCCCTGACCTTGCCCATCTGGCCGCTCCAAGTGAATGGCTGGATTTTGGCACATT
CCTGTGCAAGGTGGTCTCACTCCTGAAGGAAGTCAACTTCTATAGTGGCATCCTGCTACTGGCCTGCATC
AGTGTGGACCGTTACCTGGCCATTGTCCATGCCACACGCACACTGACCCAGAAGCGCTACTTGGTCAAAT
TCATATGTCTCAGCATCTGGGGTCTGTCCCTTGCTCCTGGCCCTGCCTGTCTACTTTCCGAAGGACCGT
CTACTCATCCAATGTTAGCCCAGCCTGCTATGAGGACATGGGCAACAATACAGCAAATGGCGGATGCTG
TTACGGATCCTGCCCCAGTCCTTTGGCTTCATCGTGCCACTGCTGATCATGCTGTTCTGTACGGATTCA
CCCTGCGTACGCTGTTTAAAGCCACATGGGGCAGAAGCACCGGGCCATGCGGGTCATCTTTGCTGTCGT
CCTCATCTTCTGCTCTGCTGGCTGCCCTACAACCTGGTCTGCTGCGCAGACACCCTCATGAGGACCCAG
GTGATCCAGGAGACCTGTGAGCGCCGAATCACATCGACCGGGCTCTGGATGCCACCGAGATTCTGGGCA
TCCTTCACAGCTGCCTCAACCCCTCATCTACGCCTTCATTGGCCAGAAGTTTCGCCATGGACTCCCTCAA
GATTCTAGCTATACATGGCTTGATCAGCAAGGACTCCCTGCCAAAGACAGCAGGCTTCCTTTGTTGGC
TCTTCTCAGGGCACACTTCCACTACTCTC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC229936 representing NM_001168298
Red=Cloning site Green=Tags(s)

MEDFNMESDSFEDFWKGEDLSNYSYSSTLPPFLDAAAPCEPESLEINKYFVVIIYALVFLLSLLGNLSLVM
LVILYSRVGRSVTDVYLLNLALADLLFALTLPIWAASKVNGWIFGTFLCKVVSLLKEVNFYSGILLLACI
SVDRYLAIHVHATRTLTKRYLVKFIKLSIWGLSLLLALPVLLFRRTVYSSNVSPACYEDMGNTANWRML
LRILPQSFQFIVPLLIMLFCYGFTRLRTLFKAHMGQKHRAMRVIFAVVLIIFLLCWLPYNLVLLADTLMTQ
VIQETCERRNHIDRALDATEILGILHSLNPLIYAFIGQKFRHGLLKILAIHGLISKDSLPKDSRPSFVG
SSSGHTSTTL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/ja1534_b09.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



* The last codon before the Stop codon of the ORF

ACCN: NM_001168298


ORF Size: 1080 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.


The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001168298.2
RefSeq ORF:	1083 bp
Locus ID:	3579
UniProt ID:	P25025
Cytogenetics:	2q35
Protein Families:	Druggable Genome, GPCR, Transmembrane
Protein Pathways:	Chemokine signaling pathway, Cytokine-cytokine receptor interaction, Endocytosis, Epithelial cell signaling in Helicobacter pylori infection
MW:	41.2 kDa
Gene Summary:	<p>The protein encoded by this gene is a member of the G-protein-coupled receptor family. This protein is a receptor for interleukin 8 (IL8). It binds to IL8 with high affinity, and transduces the signal through a G-protein activated second messenger system. This receptor also binds to chemokine (C-X-C motif) ligand 1 (CXCL1/MGSA), a protein with melanoma growth stimulating activity, and has been shown to be a major component required for serum-dependent melanoma cell growth. This receptor mediates neutrophil migration to sites of inflammation. The angiogenic effects of IL8 in intestinal microvascular endothelial cells are found to be mediated by this receptor. Knockout studies in mice suggested that this receptor controls the positioning of oligodendrocyte precursors in developing spinal cord by arresting their migration. This gene, IL8RA, a gene encoding another high affinity IL8 receptor, as well as IL8RBP, a pseudogene of IL8RB, form a gene cluster in a region mapped to chromosome 2q33-q36. Alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Nov 2009]</p>

Product images:



Circular map for RC229936



Western blot validation of overexpression lysate (Cat# [LY432936]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC229936 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).