

## **OriGene Technologies, Inc.**

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC230092L4V

## GCAT (NM\_001171690) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	GCAT (NM_001171690) Human Tagged ORF Clone Lentiviral Particle
Symbol:	GCAT
Synonyms:	KBL
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001171690
ORF Size:	1335 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC230092).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001171690.1, NP 001165161.1</u>
RefSeq ORF:	1338 bp
Locus ID:	23464
UniProt ID:	<u>075600</u>
Cytogenetics:	22q13.1
Protein Pathways:	Glycine, serine and threonine metabolism
MW:	48.4 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:The degradation of L-threonine to glycine consists of a two-step biochemical pathway<br/>involving the enzymes L-threonine dehydrogenase and 2-amino-3-ketobutyrate coenzyme A<br/>ligase. L-Threonine is first converted into 2-amino-3-ketobutyrate by L-threonine<br/>dehydrogenase. This gene encodes the second enzyme in this pathway, which then catalyzes<br/>the reaction between 2-amino-3-ketobutyrate and coenzyme A to form glycine and acetyl-<br/>CoA. The encoded enzyme is considered a class II pyridoxal-phosphate-dependent<br/>aminotransferase. Alternate splicing results in multiple transcript variants. A pseudogene of<br/>this gene is found on chromosome 14. [provided by RefSeq, Jan 2010]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US