

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for RC230191L1V

## HCK (NM\_001172131) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	HCK (NM_001172131) Human Tagged ORF Clone Lentiviral Particle
Symbol:	НСК
Synonyms:	JTK9; p59Hck; p61Hck
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_001172131
ORF Size:	1512 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC230191).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001172131.1</u>
RefSeq ORF:	1515 bp
Locus ID:	3055
UniProt ID:	<u>P08631</u>
Cytogenetics:	20q11.21
Protein Families:	Druggable Genome, Protein Kinase
Protein Pathways:	Chemokine signaling pathway, Fc gamma R-mediated phagocytosis
MW:	57.7 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



#### HCK (NM\_001172131) Human Tagged ORF Clone Lentiviral Particle – RC230191L1V

Gene Summary:The protein encoded by this gene is a member of the Src family of tyrosine kinases. This<br/>protein is primarily hemopoietic, particularly in cells of the myeloid and B-lymphoid lineages.<br/>It may help couple the Fc receptor to the activation of the respiratory burst. In addition, it may<br/>play a role in neutrophil migration and in the degranulation of neutrophils. Multiple isoforms<br/>with different subcellular distributions are produced due to both alternative splicing and the<br/>use of alternative translation initiation codons, including a non-AUG (CUG) codon. [provided<br/>by RefSeq, Feb 2010]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US