

Product datasheet for RC230417L3

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

splicing factor 1 (SF1) (NM_001178030) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: splicing factor 1 (SF1) (NM_001178030) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: splicing factor 1

Synonyms: BBP; D11S636; MBBP; ZCCHC25; ZFM1; ZNF162

Mammalian Cell Pu

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)


E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC230417).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:






^{*} The last codon before the Stop codon of the ORF.





Plasmid Map:



ACCN: NM_001178030

ORF Size: 2019 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

> reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

NM 001178030.1 RefSeq:

RefSeq ORF: 2022 bp

Locus ID: 7536





UniProt ID: Q15637
Cytogenetics: 11q13.1

Protein Families: Transcription Factors

MW: 72.2 kDa

Gene Summary: This gene encodes a nuclear pre-mRNA splicing factor. The encoded protein specifically

recognizes the intron branch point sequence at the 3' splice site, together with the large subunit of U2 auxiliary factor (U2AF), and is required for the early stages of spliceosome assembly. It also plays a role in nuclear pre-mRNA retention and transcriptional repression. The encoded protein contains an N-terminal U2AF ligand motif, a central hnRNP K homology motif and quaking 2 region which bind a key branch-site adenosine within the branch point sequence, a zinc knuckles domain, and a C-terminal proline-rich domain. Alternative splicing

results in multiple transcript variants. [provided by RefSeq, Oct 2016]