

## Product datasheet for RC230616L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MAN2B1 (NM\_001173498) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** MAN2B1 (NM\_001173498) Human Tagged ORF Clone Lentiviral Particle

Symbol: MAN2B1

Synonyms: LAMAN; MANB

Mammalian Cell

viaitiitialiati Celi

Puromycin

Selection: Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001173498

ORF Size: 3030 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC230616).

Sequence:

Cytogenetics:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** NM 001173498.1, NP 001166969.1

 RefSeq Size:
 3239 bp

 RefSeq ORF:
 3033 bp

 Locus ID:
 4125

 UniProt ID:
 000754

**Protein Families:** Druggable Genome

**Protein Pathways:** Lysosome, Other glycan degradation

19p13.13





ORIGENE

**MW:** 113.6 kDa

**Gene Summary:** 

This gene encodes an enzyme that hydrolyzes terminal, non-reducing alpha-D-mannose residues in alpha-D-mannosides. Its activity is necessary for the catabolism of N-linked carbohydrates released during glycoprotein turnover and it is member of family 38 of glycosyl hydrolases. The full length protein is processed in two steps. First, a 49 aa leader sequence is cleaved off and the remainder of the protein is processed into 3 peptides of 70 kDa, 42 kDa (D) and 13/15 kDa (E). Next, the 70 kDa peptide is further processed into three peptides (A, B and C). The A, B and C peptides are disulfide-linked. Defects in this gene have been associated with lysosomal alpha-mannosidosis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Mar 2010]