

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC230863L4

DDX3 (DDX3X) (NM_001193416) Human Tagged Lenti ORF Clone

Product data:

| Product Type: | Expression Plasmids | | |
|------------------------------|---|--|--|
| Product Name: | DDX3 (DDX3X) (NM_001193416) Human Tagged Lenti ORF Clone | | |
| Tag: | mGFP | | |
| Symbol: | DDX3 | | |
| Synonyms: | CAP-Rf; DBX; DDX3; DDX14; HLP2; MRX102; MRXSSB | | |
| Mammalian Cell Selection: | Puromycin | | |
| Vector: | pLenti-C-mGFP-P2A-Puro (PS100093) | | |
| E. coli Selection: | Chloramphenicol (34 ug/mL) | | |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(RC230863). | | |
| Restriction Sites: | Sgfl-Mlul | | |
| Cloning Scheme: | | | |
| | Cloning sites used for ORF Shuttling: | | |
| | Sgf I ORF Mlu I GCG ATC GCC ATG NNN ACG CGT | | |




* The last codon before the Stop codon of the ORF.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Plasmid Map:



| ACCN: | NM_001193416 |
|------------------------|--|
| ORF Size: | 1983 bp |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u> |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | <u>NM 001193416.2</u> |
| RefSeq Size: | 4662 bp |
| RefSeq ORF: | 1986 bp |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

| | ~ | ×″/ | - | |
|--------|-----------|-----|-----|----|
| | () | 2ÎC | .FN | VF |
| \sim | | ~ | | |

| Locus ID: UniProt ID: | 1654 <u>000571</u> |
|--------------------------|--|
| Cytogenetics: | Xp11.4 |
| Protein Families: | ES Cell Differentiation/IPS |
| Protein Pathways: | RIG-I-like receptor signaling pathway |
| MW: | 73.2 kDa |
| Gene Summary: | The protein encoded by this gene is a member of the large DEAD-box protein family, that is defined by the presence of the conserved Asp-Glu-Ala-Asp (DEAD) motif, and has ATP-dependent RNA helicase activity. This protein has been reported to display a high level of RNA-independent ATPase activity, and unlike most DEAD-box helicases, the ATPase activity is thought to be stimulated by both RNA and DNA. This protein has multiple conserved domains and is thought to play roles in both the nucleus and cytoplasm. Nuclear roles include transcriptional regulation, mRNP assembly, pre-mRNA splicing, and mRNA export. In the cytoplasm, this protein is thought to be involved in translation, cellular signaling, and viral replication. Misregulation of this gene has been implicated in tumorigenesis. This gene has a paralog located in the nonrecombining region of the Y chromosome. Pseudogenes sharing similarity to both this gene and the DDX3Y paralog are found on chromosome 4 and the X chromosome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2014] |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US