

## Product datasheet for RC230968L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MD2 (LY96) (NM\_001195797) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** MD2 (LY96) (NM\_001195797) Human Tagged ORF Clone Lentiviral Particle

Symbol: MD2

**Synonyms:** ESOP-1; ly-96; MD-2; MD2

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001195797

ORF Size: 390 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC230968).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** NM 001195797.1, NP 001182726.1

 RefSeq Size:
 552 bp

 RefSeq ORF:
 393 bp

 Locus ID:
 23643

 UniProt ID:
 Q9Y6Y9

Cytogenetics: 8q21.11

**Protein Families:** Secreted Protein

**Protein Pathways:** Pathogenic Escherichia coli infection, Toll-like receptor signaling pathway





ORIGENE

**MW:** 14.9 kDa

**Gene Summary:** This gene encodes a protein which associates with toll-like receptor 4 on the cell surface and

confers responsiveness to lipopolysaccyaride (LPS), thus providing a link between the receptor and LPS signaling. Studies of the mouse ortholog suggest that this gene may be involved in endotoxin neutralization. Alternative splicing results in multiple transcript variants

encoding different isoforms. [provided by RefSeq, Sep 2010]