

# Product datasheet for RC230991

## ATP6V1F (NM\_001198909) Human Tagged ORF Clone

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


Product Type:	Expression Plasmids
Product Name:	ATP6V1F (NM_001198909) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	ATP6V1F
Synonyms:	ATP6S14; VATF; Vma7
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin
ORF Nucleotide Sequence:	>RC230991 representing NM_001198909 Red=Cloning site Blue=ORF Green=Tags(s)
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGGCGGGGAGGGTAAGCTCATCGCAGTGATCGGAGACGAGGACACGGTGACTGGTTTCCTGCTGGGCG GCATAGGGGAGCTTAACAAGAACCGCCATCCCAATTTCCTGGTGGTGGAGAAGGATACAACCATCAATGA GATCGAAGACACTTTCCGTTCACTTGGAAGCCTTCCGGGCAGTGTTGTAGAAGCCAACCCTAATCAGCGT GACCCTCCGCTTTGGGATGAAATTGATTCTAGGCAATTTCTAAACCGGGATGACATTGGCATCATCCTCA TCAACCAGTACATCGCAGAGATGGTGCGGCATGCCCTGGACGCCCACCAGCAGTCCATCCCCGCTGTCCT GGAGATCCCCTCCAAGGAGCACCCATATGACGCCGCCAAGGACTCCATCCTGCGCAGGGCCAGGGGCATG TTCACTGCCGAAGACCTGCGC
	ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAG <b>GTTTAA</b>
Protein Sequence:	>RC230991 representing NM_001198909 <mark>Red</mark> =Cloning site Green=Tags(s)
	MAGRGKLIAVIGDEDTVTGFLLGGIGELNKNRHPNFLVVEKDTTINEIEDTFRSLGSLPGSVVEANPNQR DPPLWDEIDSRQFLNRDDIGIILINQYIAEMVRHALDAHQQSIPAVLEIPSKEHPYDAAKDSILRRARGM FTAEDLR
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
<b>Restriction Sites:</b>	Sgfl-Mlul



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US




#### **Cloning Scheme:**



\* The last codon before the Stop codon of the ORF

### Plasmid Map:



ACCN:	NM_001198909
ORF Size:	441 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CRIGENE ATP6V1F (NM_001198909) Human Tagged ORF Clone – RC230991	
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
RefSeq:	<u>NM 001198909.2</u>
RefSeq ORF:	444 bp
Locus ID:	9296
UniProt ID:	<u>Q16864</u>
Cytogenetics:	7q32.1
Protein Pathways:	Epithelial cell signaling in Helicobacter pylori infection, Metabolic pathways, Oxidative phosphorylation, Vibrio cholerae infection
MW:	16.9 kDa
Gene Summary:	This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c", and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is the V1 domain F subunit protein. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US