

## Product datasheet for RC231053L4

### glutathione S transferase Omega 1 (GSTO1) (NM\_001191002) Human Tagged Lenti ORF Clone

#### Product data:

|                           |   |
|---------------------------|---|
| Product Type:             | Expression Plasmids   |
| Product Name:             | glutathione S transferase Omega 1 (GSTO1) (NM_001191002) Human Tagged Lenti ORF Clone |
| Tag:                      | mGFP  |
| Symbol:                   | glutathione S transferase Omega 1   |
| Synonyms:                 | GSTO 1-1; GSTTLp28; HEL-S-21; P28; SPG-R  |
| Mammalian Cell Selection: | Puromycin   |
| Vector:                   | pLenti-C-mGFP-P2A-Puro (PS100093)   |
| E. coli Selection:        | Chloramphenicol (34 ug/mL)  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC231053).                        |
| Restriction Sites:        | SgfI-RsrII  |
| Cloning Scheme:           |   |

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


|           |              |
|-----------|--------------|
| ACCN:     | NM_001191002 |
| ORF Size: | 624 bp       |



[View online >](#)

|                               |   |
|-------------------------------|---|
| <b>OTI Disclaimer:</b>        | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>  |
| <b>OTI Annotation:</b>        | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.  |
| <b>Components:</b>            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).  |
| <b>Reconstitution Method:</b> | <ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol> |
| <b>RefSeq:</b>                | <a href="#">NM_001191002.1</a> , <a href="#">NP_001177931.1</a>   |
| <b>RefSeq ORF:</b>            | 627 bp  |
| <b>Locus ID:</b>              | 9446  |
| <b>UniProt ID:</b>            | <a href="#">P78417</a>  |
| <b>Cytogenetics:</b>          | 10q25.1   |
| <b>Protein Families:</b>      | Druggable Genome  |
| <b>Protein Pathways:</b>      | Drug metabolism - cytochrome P450, Glutathione metabolism, Metabolism of xenobiotics by cytochrome P450   |
| <b>MW:</b>                    | 24.2 kDa  |
| <b>Gene Summary:</b>          | The protein encoded by this gene is an omega class glutathione S-transferase (GST) with glutathione-dependent thiol transferase and dehydroascorbate reductase activities. GSTs are involved in the metabolism of xenobiotics and carcinogens. The encoded protein acts as a homodimer and is found in the cytoplasm. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2010]   |

Product images:



Circular map for RC231053L4