

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC231175L3V


Caspase 12 (CASP12) (NM_001191016) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Caspase 12 (CASP12) (NM_001191016) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Caspase 12
Synonyms:	CASP-12; CASP12P1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001191016
ORF Size:	1023 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC231175).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001191016.1, NP 001177945.1</u>
RefSeq ORF:	1026 bp
Locus ID:	100506742
UniProt ID:	Q6UXS9
Cytogenetics:	11q22.3
MW:	39.4 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Caspases are cysteine proteases that cleave C-terminal aspartic acid residues on their
substrate molecules. This gene is most highly related to members of the ICE subfamily of
caspases that process inflammatory cytokines. In rodents, the homolog of this gene mediates
apoptosis in response to endoplasmic reticulum stress. However, in humans this gene
contains a polymorphism for the presence or absence of a premature stop codon. The
majority of human individuals have the premature stop codon and produce a truncated non-
functional protein. The read-through codon occurs primarily in individuals of African descent
and carriers have endotoxin hypo-responsiveness and an increased susceptibility to severe
sepsis. Several alternatively spliced transcript variants have been noted for this gene.
[provided by RefSeq, Feb 2011]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US