

Product datasheet for RC231175L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Caspase 12 (CASP12) (NM_001191016) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Caspase 12 (CASP12) (NM 001191016) Human Tagged ORF Clone Lentiviral Particle

Symbol: Caspase 12

Synonyms: CASP-12; CASP12P1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001191016

ORF Size: 1023 bp

ORF Nucleotide

_.

Sequence:

UniProt ID:

The ORF insert of this clone is exactly the same as(RC231175).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001191016.1, NP 001177945.1

Q6UXS9

RefSeq ORF: 1026 bp

Locus ID: 100506742

Cytogenetics: 11q22.3

MW: 39.4 kDa





Caspase 12 (CASP12) (NM_001191016) Human Tagged ORF Clone Lentiviral Particle – RC231175L4V

Gene Summary:

Caspases are cysteine proteases that cleave C-terminal aspartic acid residues on their substrate molecules. This gene is most highly related to members of the ICE subfamily of caspases that process inflammatory cytokines. In rodents, the homolog of this gene mediates apoptosis in response to endoplasmic reticulum stress. However, in humans this gene contains a polymorphism for the presence or absence of a premature stop codon. The majority of human individuals have the premature stop codon and produce a truncated nonfunctional protein. The read-through codon occurs primarily in individuals of African descent and carriers have endotoxin hypo-responsiveness and an increased susceptibility to severe sepsis. Several alternatively spliced transcript variants have been noted for this gene. [provided by RefSeq, Feb 2011]