

## Product datasheet for RC231201L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cytochrome P450 2C8 (CYP2C8) (NM\_001198854) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Cytochrome P450 2C8 (CYP2C8) (NM\_001198854) Human Tagged ORF Clone Lentiviral Particle

Symbol: Cytochrome P450 2C8

Synonyms: CPC8; CYP2C8DM; CYPIIC8; MP-12/MP-20

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_001198854

ORF Size: 1164 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC231201).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 001198854.1, NP 001185783.1</u>

 RefSeq ORF:
 1167 bp

 Locus ID:
 1558

 UniProt ID:
 P10632

 Cytogenetics:
 10q23.33

**Protein Families:** Druggable Genome, P450, Transmembrane

**Protein Pathways:** Arachidonic acid metabolism, Drug metabolism - cytochrome P450, Linoleic acid metabolism,

Metabolic pathways, Metabolism of xenobiotics by cytochrome P450, Retinol metabolism





MW: 44.8 kDa

**Gene Summary:** This gene encodes a member of the cytochrome P450 superfamily of enzymes. The

cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by phenobarbital. The enzyme is known to metabolize many xenobiotics, including the anticonvulsive drug mephenytoin, benzo(a)pyrene, 7-ethyoxycoumarin, and the anti-cancer drug taxol. This gene is located within a cluster of cytochrome P450 genes on chromosome 10q24. Several transcript variants encoding a few different isoforms have been found for this gene. [provided by RefSeq, Nov

2010]