

## Product datasheet for RC231386L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PDE1C (NM\_001191057) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** PDE1C (NM\_001191057) Human Tagged ORF Clone Lentiviral Particle

Symbol: PDE10

**Synonyms:** cam-PDE 1C; DFNA74; hCam-3; Hcam3

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001191057

ORF Size: 2127 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC231386).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. <u>More info</u>

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 001191057.1</u>

**RefSeq ORF:** 2130 bp **Locus ID:** 5137

UniProt ID: Q14123
Cytogenetics: 7p14.3

**Protein Families:** Druggable Genome

**Protein Pathways:** Calcium signaling pathway, Olfactory transduction, Progesterone-mediated oocyte

maturation, Purine metabolism







MW:

81.2 kDa

**Gene Summary:** 

This gene encodes an enzyme that belongs to the 3'5'-cyclic nucleotide phosphodiesterase family. Members of this family catalyze hydrolysis of the cyclic nucleotides, cyclic adenosine monophosphate and cyclic guanosine monophosphate, to the corresponding nucleoside 5'-monophosphates. The enzyme encoded by this gene regulates proliferation and migration of vascular smooth muscle cells, and neointimal hyperplasia. This enzyme also plays a role in pathological vascular remodeling by regulating the stability of growth factor receptors, such as PDGF-receptor-beta. [provided by RefSeq, Jul 2016]