

## Product datasheet for **RC231398L3V**

### **PDE4D (NM\_001197218) Human Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | PDE4D (NM_001197218) Human Tagged ORF Clone Lentiviral Particle  |
| Symbol:                   | PDE4D  |
| Synonyms:                 | ACRDYS2; DPDE3; HSPDE4D; PDE4DN2; PDE43; STRK1   |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-Myc-DDK-P2A-Puro (PS100092)   |
| Tag:                      | Myc-DDK  |
| ACCN:                     | NM_001197218   |
| ORF Size:                 | 2235 bp  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC231398).   |
| OTI Disclaimer:           | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">NM_001197218.1</a>   |
| RefSeq ORF:               | 2238 bp  |
| Locus ID:                 | 5144   |
| UniProt ID:               | <a href="#">Q08499</a>   |
| Cytogenetics:             | 5q11.2-q12.1   |
| Protein Families:         | Druggable Genome   |
| Protein Pathways:         | Progesterone-mediated oocyte maturation, Purine metabolism   |
| MW:                       | 84.9 kDa   |



[View online »](#)

**Gene Summary:**

This gene encodes one of four mammalian counterparts to the fruit fly 'dunce' gene. The encoded protein has 3',5'-cyclic-AMP phosphodiesterase activity and degrades cAMP, which acts as a signal transduction molecule in multiple cell types. This gene uses different promoters to generate multiple alternatively spliced transcript variants that encode functional proteins.[provided by RefSeq, Sep 2009]