

Product datasheet for RC231440L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

NCoR (NCOR1) (NM_001190438) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Symbol: NCoR

Synonyms: hN-CoR; N-CoR; N-CoR1; PPP1R109; TRAC1

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001190438

ORF Size: 2742 bp

ORF Nucleotide Sequence: The ORF insert of this clone is exactly the same as(RC231440).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM_001190438.1</u>, <u>NP_001177367.1</u>

RefSeq Size: 3106 bp

RefSeq ORF: 2745 bp

Locus ID: 9611

UniProt ID: <u>075376</u>

Cytogenetics: 17p12-p11.2



NCoR (NCOR1) (NM_001190438) Human Tagged ORF Clone Lentiviral Particle | RC231440L3V

Protein Families: Druggable Genome

MW: 104.4 kDa

Gene Summary: This gene encodes a protein that mediates ligand-independent transcription repression of

thyroid-hormone and retinoic-acid receptors by promoting chromatin condensation and preventing access of the transcription machinery. It is part of a complex which also includes histone deacetylases and transcriptional regulators similar to the yeast protein Sin3p. This gene is located between the Charcot-Marie-Tooth and Smith-Magenis syndrome critical regions on chromosome 17. Alternate splicing results in multiple transcript variants.

Pseudogenes of this gene are found on chromosomes 17 and 20.[provided by RefSeq, Jun

2010