

Product datasheet for RC231501L3

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PEG10 (NM_001184961) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: PEG10 (NM 001184961) Human Tagged Lenti ORF Clone

Tag: Myc-DDK
Symbol: PEG10

Synonyms: EDR; HB-1; Mar2; Mart2; MEF3L; RGAG3; RTL2; SIRH1

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)


E. coli Selection: Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(RC231501).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_001184961

ORF Size: 2066 bp



PEG10 (NM_001184961) Human Tagged Lenti ORF Clone - RC231501L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001184961.1</u>

 RefSeq ORF:
 2228 bp

 Locus ID:
 23089

 UniProt ID:
 Q86TG7

 Cytogenetics:
 7q21.3

 MW:
 85.05 kDa


Gene Summary: This is a paternally expressed imprinted gene that is thought to have been derived from the

Ty3/Gypsy family of retrotransposons. It contains two overlapping open reading frames, RF1 and RF2, and expresses two proteins: a shorter, gag-like protein (with a CCHC-type zinc finger domain) from RF1; and a longer, gag/pol-like fusion protein (with an additional aspartic protease motif) from RF1/RF2 by -1 translational frameshifting (-1 FS). While -1 FS has been observed in RNA viruses and transposons in both prokaryotes and eukaryotes, this gene represents the first example of -1 FS in a eukaryotic cellular gene. This gene is highly conserved across mammalian species and retains the heptanucleotide (GGGAAAC) and pseudoknot elements required for -1 FS. It is expressed in adult and embryonic tissues (most notably in placenta) and reported to have a role in cell proliferation, differentiation and apoptosis. Overexpression of this gene has been associated with several malignancies, such as hepatocellular carcinoma and B-cell lymphocytic leukemia. Knockout mice lacking this gene showed early embryonic lethality with placental defects, indicating the importance of this gene in embryonic development. Additional isoforms resulting from alternatively spliced transcript variants, and use of upstream non-AUG (CUG) start codon have been reported for

this gene. [provided by RefSeq, Oct 2014]



Product images:



Circular map for RC231501L3