

Product datasheet for RC233265

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

Insulin (INS) (NM 001185097) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Insulin (INS) (NM_001185097) Human Tagged ORF Clone

Tag: Myc-DDK Symbol: Insulin

Synonyms: IDDM; IDDM1; IDDM2; ILPR; IRDN; MODY10; PNDM4

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-Entry (PS100001) E. coli Selection: Kanamycin (25 ug/mL) >RC233265 ORF sequence **ORF Nucleotide**

Red=Cloning site Blue=ORF Green=Tags(s) Sequence:

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGCCCTGTGGATGCGCCTCCTGCCCCTGCTGGCGCTGCTGGCCCTCTGGGGACCTGACCCAGCCGCAG CCTTTGTGAACCAACACCTGTGCGGCTCACACCTGGTGGAAGCTCTCTACCTAGTGTGCGGGGAACGAGG CTTCTTCTACACCCCAAGACCCGCGGGAGGCAGAGGACCTGCAGGTGGGGCAGGTGGAGCTGGGCGGG GGCCCTGGTGCAGGCAGCCTTGCAGCCCTTGGCGGAGGGGTCCCTGCAGAAGCGTGGCATTGTGGAAC AATGCTGTACCAGCATCTGCTCCCTCTACCAGCTGGAGAACTACTGCAAC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

>RC233265 protein sequence **Protein Sequence:**

Red=Cloning site Green=Tags(s)

MALWMRLLPLLALLALWGPDPAAAFVNQHLCGSHLVEALYLVCGERGFFYTPKTRREAEDLQVGQVELGG

GPGAGSLQPLALEGSLQKRGIVEQCCTSICSLYQLENYCN

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6050_e08.zip


Restriction Sites: Sgfl-Mlul





Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_001185097

ORF Size: 330 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 001185097.2

RefSeq Size: 495 bp RefSeq ORF: 333 bp Locus ID: 3630



UniProt ID: P01308 Cytogenetics: 11p15.5

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Secreted Protein

Insulin signaling pathway, Maturity onset diabetes of the young, mTOR signaling pathway, **Protein Pathways:**

Oocyte meiosis, Progesterone-mediated oocyte maturation, Prostate cancer, Regulation of


actin cytoskeleton, Regulation of autophagy, Type I diabetes mellitus, Type II diabetes mellitus

MW: 12 kDa

Gene Summary: This gene encodes insulin, a peptide hormone that plays a vital role in the regulation of

carbohydrate and lipid metabolism. After removal of the precursor signal peptide, proinsulin is post-translationally cleaved into three peptides: the B chain and A chain peptides, which are covalently linked via two disulfide bonds to form insulin, and C-peptide. Binding of insulin to the insulin receptor (INSR) stimulates glucose uptake. A multitude of mutant alleles with phenotypic effects have been identified, including insulin-dependent diabetes mellitus, permanent neonatal diabetes diabetes mellitus, maturity-onset diabetes of the young type 10 and hyperproinsulinemia. There is a read-through gene, INS-IGF2, which overlaps with this gene at the 5' region and with the IGF2 gene at the 3' region. [provided by RefSeq, May 2020]

Product images:



Circular map for RC233265