

Product datasheet for **RC233535**

UCP4 (SLC25A27) (NM_001204052) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	UCP4 (SLC25A27) (NM_001204052) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	UCP4
Synonyms:	UCP4
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC233535 representing NM_001204052 Red=Cloning site Blue=ORF Green=Tags(s)

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGTCCGTCCCGAGGAGGAGGAGAGGCTTTTGCCGTGACCCAGAGATGGCCCCGAGCGAGCAAATTC
 TACTGTCCGGCTGCGCGGCTACCGTGGCCGAGCTAGCAACCTTTCCCTGGATCTCACAAAACCTCGACT
 CCAAATGCAAGGAGAAGCAGCTCTTGCTCGTTGGGAGACGGTCAAGAGAATCTGCCCTATAGGGGA
 ATGGTGCACAGCTCTAGGGATCATTGAAGAGGAAGGCTTTCTAAAGCTTTGGCAAGGAGTGACACCCG
 CCATTTACAGACACGTAGTGTATTCTGGAGGTCAATGGTCACATATGAACATCTCCGAGAGGTTGTGTT
 TGGCAAAAGTGAAGATGAGCATTATCCCTTTGGAAATCAGTCATTGGAGGGATGATGGCTGGTGTATT
 GGCCAGTTTTAGCCAATCCAACCTGACCTAGTGAAGGTTCAAGTGAATGGAAGGAAAAAGGAACTGG
 AAGGAAAACCATTCGATTTCTGGGTGTACATCATGCAATTTGCAAAAATCTTAGCTGAAGGAGGAATACG
 AGGGCTTTGGGCAGGCTGGGTACCAATATACAAAGAGCAGCACTGGTGAATATGGGAGATTTAACCCT
 TATGATACAGTGAACACTACTTGGTATTGAATACCACTTGAGGACAATATCATGACTCACGGTTTAT
 CAAGTGATCTGGTCGGATCTCACAAGGCCATCCAA

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA


[View online »](#)


Protein Sequence: >RC233535 representing NM_001204052
 Red=Cloning site Green=Tags(s)

MSVPEEEERLLPLTQRWPRASKFLLSGCAATVAELATFPLDLTKTRLQMGEAALARLGDGARESAPYRG
 MVRTALGIIIEEGFLKLWQGVTPAIYRHVVYSGGRMVTYEHLREVVFGESEDEHYPLWKSIVGGMMAGVI
 GQFLANPTDLVKVQMMEGKRKLEGGKPLRFRGVHHAFAKILAEAGGIRGLWAGWVPNIQRAALVNMGDLTT
 YDTVKHLYLVNTPLEDNIMTHGLSSDLVGSKAIQ

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001204052

ORF Size: 735 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM_001204052.1, NP_001190981.1</u>
RefSeq Size:	2526 bp
RefSeq ORF:	738 bp
Locus ID:	9481
UniProt ID:	<u>O95847</u>
Cytogenetics:	6p12.3
Protein Families:	Druggable Genome
MW:	27.7 kDa
Gene Summary:	<p>Mitochondrial uncoupling proteins (UCP) are members of the larger family of mitochondrial anion carrier proteins (MACP). UCPs separate oxidative phosphorylation from ATP synthesis with energy dissipated as heat, also referred to as the mitochondrial proton leak. UCPs facilitate the transfer of anions from the inner to the outer mitochondrial membrane and the return transfer of protons from the outer to the inner mitochondrial membrane. They also reduce the mitochondrial membrane potential in mammalian cells. Tissue specificity occurs for the different UCPs and the exact methods of how UCPs transfer H⁺/OH⁻ are not known. UCPs contain the three homologous protein domains of MACPs. Transcripts of this gene are only detected in brain tissue and are specifically modulated by various environmental conditions. Alternative splicing results in multiple transcript variants.[provided by RefSeq, Feb 2011]</p>

Product images:



Circular map for RC233535