

## **Product datasheet for RC233597**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Prolactin Receptor (PRLR) (NM\_001204318) Human Tagged ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Prolactin Receptor (PRLR) (NM\_001204318) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: Prolactin Receptor

Synonyms: HPRL; hPRLrl; MFAB; RI-PRLR

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >RC233597 representing NM\_001204318
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA





Protein Sequence: >RC233597 representing NM\_001204318

Red=Cloning site Green=Tags(s)


MKENVASATVFTLLLFLNTCLLNGQLPPGKPEIFKCRSPNKETFTCWWRPGTDGGLPTNYSLTYHREGET LMHECPDYITGGPNSCHFGKQYTSMWRTYIMMVNATNQMGSSFSDELYVDVTYIVQPDPPLELAVEVKQP EDRKPYLWIKWSPPTLIDLKTGWFTLLYEIRLKPEKAAEWEIHFAGQQTEFKILSLHPGQKYLVQVRCKP DHGYWSAWSPATFIQIPSGDPLMLGASHYKNLKSYRPRKISSQGRLAVFTKATLTTVQ


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** 

Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_001204318

ORF Size: 804 bp

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customercom">customercom</a> or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.



#### Prolactin Receptor (PRLR) (NM\_001204318) Human Tagged ORF Clone - RC233597

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001204318.1</u>, <u>NP 001191247.1</u>

RefSeq Size: 1461 bp
RefSeq ORF: 807 bp
Locus ID: 5618
UniProt ID: P16471
Cytogenetics: 5p13.2

**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway, Neuroactive ligand-

receptor interaction

MW: 31.2 kDa


**Gene Summary:** This gene encodes a receptor for the anterior pituitary hormone, prolactin, and belongs to

the type I cytokine receptor family. Prolactin-dependent signaling occurs as the result of ligand-induced dimerization of the prolactin receptor. Several alternatively spliced transcript variants encoding different membrane-bound and soluble isoforms have been described for this gene, which may function to modulate the endocrine and autocrine effects of prolactin in

normal tissue and cancer. [provided by RefSeq, Feb 2011]



# **Product images:**



Circular map for RC233597