

Product datasheet for **RC234672**

DAXX (NM_001254717) Human Tagged ORF Clone

Product data:

| | |
|---------------------------|--|
| Product Type: | Expression Plasmids |
| Product Name: | DAXX (NM_001254717) Human Tagged ORF Clone |
| Tag: | Myc-DDK |
| Symbol: | DAXX |
| Synonyms: | BING2; DAP6; EAP1; SMIM40 |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |



[View online »](#)

ORF Nucleotide Sequence:

>RC234672 representing NM_001254717
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGCAGACAGCAGACCACCTGAGTGGTCCCATTCTCTATAACCGGCAGCAACGTGCCACTCTCTGT
 TTTTGGCCTCGGCGGAGTTCTGCAACATCCTCTCTAGGGTCTGTCTCGGCCCGGAGCCGGCCAGCCAA
 GCTCTATGTCTACATCAATGAGCTCTGCACTGTTCTCAAGGCCCACTCAGCCAAAAAGAAGCTGAACTTG
 GCCCTGCCGCCACCACCTCCAATGAGCCCTCTGGGAATAAACCTCCCACACACCTCTCCTTGGACCCCA
 CAAATGCTGAAAACACTGCCTCTCAGTCTCAAGGACCGTGGTCCCGGCGCAGATCCAGCGTTTGA
 GCAGCTGTGGCCTCTATGTGGCAGAGATCCGGCGGTGCAGGAAAAGGAGTTGGATCTCTCAGAATTG
 GATGACCCAGACTCCGCATACCTGCAGGAGGCACGGTTGAAGCGTAAGCTGATCCGCCTCTTTGGGCGAC
 TATGTGAGCTGAAAGACTGCTCTTCACTGACCGCCGTGTATAGAGCAGCGCATCCCCTACCGTGGCAC
 CCGCTACCCAGAGGTTAACAGGGCATTGAGCGGCTCATCAACAAGCCAGGGCCTGATACCTTCCCTGAC
 TATGGGGATGTGCTTCGGGCTGTAGAGAAGGCAGCTGCCCGACACAGCCTTGGCCTCCCCGACAGCAGC
 TCCAGCTCATGGCTCAGGATGCCTCCGAGATGTGGGCATCAGGTTACAGGAGCGACGTCACCTCGATCT
 CATCTACAACCTTTGGCTGCCACCTCACAGATGACTATAGGCCAGGCGTTGACCCTGCACTATCAGATCCT
 GTGTTGGCCCGCGCCTTCGGGAAAACCGGAGTTTGGCCATGAGTCGGCTGGATGAGGTCATCTCCAAAT
 ATGCAATGTTGCAAGACAAAAGTGAAGGAGGCGAGAGAAAAAGAGAAGAGCTCGGCTCCAAGGCACCTC
 TCCCCTCTGCAGACACCCCGAAGCCTCCTTGGATTCTGGTGAAGGCGCCTAGTGAATGGCATCCCAG
 GGGTGCCTTCTGCCTCCAGAGCTGAGACAGATGACGAAGACGATGAGGAGAGTGATGAGGAAGAGGAGG
 AGGAGGAGGAAGAAGAAGAGGAGGAGGCCACAGATTCTGAAGAGGAGGAGGATCTGGAACAGATGCAGGA
 GGGTCAGGAGGATGATGAAGAGGAGGACGAAGAGGAAGAAGCAGCAGGTAAGATGGAGACAAGAGC
 CCCATGTCCTCACTACAGATCTCCAATGAAAAGAACCTGGAACCTGGCAAACAGATCAGCAGATCTTCAG
 GGGAGCAGCAAAACAAAGGACGCATAGTGTACCATCGTTACTGTGAGAAGAACCCTGGCCCCCTCCAG
 CATAGATGCTGAAAGCAATGGAGAACAGCCTGAGGAGCTGACCCTGGAGGAAGAAAGCCCTGTGTCTCAG
 CTCTTTGAGCTAGAGATTGAAGCTTTGCCCTGGATACCCCTTCTCTGTGGAGACGGACATTTCTCTTT
 CCAGGAAGCAATCAGAGGAGCCCTTCACTGTCTTAGAGAATGGAGCAGGCATGGTCTCTTCTACTTC
 CTTCAATGGAGGCGTCTCTCTCACAACGGGGAGATTCTGGTCCCCCTGCAAAAAATCTCGGAAGGAG
 AAGAAGCAACAGGATCAGGGCCATTAGGAAACAGCTATGTGAAAGGCAAAGGTCAGTGCATGAGAAGA
 ATGGGAAAAAGATATGTACCCTGCCAGCCACCTTCCCCTTGGCTTCTTGGCCCCAGTTGCTGATTC
 CTCCACGAGGGTGGACTCTCCAGCCATGGCCTGGTGACCAGCTCCCTCTGCATCCCTTCTCCAGCCCGG
 CTGTCCAAACCCCCATTACAGCCTCCTCGGCCTGGTACTTGAAGACAAGTGTGGCCACACAATGCG
 ATCCAGAAGAGATCATCGTGCTCTCAGACTCTGAT

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC234672 representing NM_001254717
Red=Cloning site Green=Tags(s)

MQTADHPEVVPFLYNRQQRAHSLFLASAEFCNILSRVLSRARSRAKLYVYINELCTVLKAHSAKKNLNL
 APAATTSNEPSGNNPPTHLSLDPTNAENTASQSPRTRGSRRQIQRLQLLALYVAEIRRLQEKLDLSEL
 DDPDSAYLQEARLKRKLRIRLFGRLCELKDCSSLTGRVIEQRIPIYRGTRYPEVNRRIERL INKPGPDTFPD
 YGDVLRAVEKAAARHSLGLPRQQLQLMAQDAFRDVGIRLQERRHLDLIYNFGCHL TDDYRPGVDPALSDP
 VLARRLRENRLAMSRLDEVISKYAMLQDKSEEGERKKRRARLQGTSSHSADTPEASLDSGEGPSGMASQ
 GCPASRAETDDEDEESDEEEEEEEEEEEEAATDSEEEEDLEQMMEGQEDDEEEDEEEAAAGKDGDKS
 PMSSLQISNEKNLEPGKQISRSSGEQNKGRIVSPSLLSEEPLAPSSIDAESNGEQPEEL TLEESPVSQ
 LFELEIEALPLDTPSSVETDISSSRKQSEEPFTTVLENGAGMVSSTSFNGGVSPHNWGDSPGPKKSRKE
 KKQTGSGPLGNSYVERQSVHEKNGKIKTLPSPPLASLAPVADSSTRVDSPSHGLVTSLLCIPSPAR
 LSQTPHSQPPRPGTCKTSVATQCDPEEIIVLSDSD

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:



ACCN: NM_001254717

ORF Size: 1995 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

| | |
|-------------------------------|---|
| Reconstitution Method: | <ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | NM_001254717.1 , NP_001241646.1 |
| RefSeq Size: | 2369 bp |
| RefSeq ORF: | 1998 bp |
| Locus ID: | 1616 |
| UniProt ID: | Q9UER7 |
| Cytogenetics: | 6p21.32 |
| Protein Families: | Druggable Genome, Stem cell - Pluripotency, Transcription Factors |
| Protein Pathways: | Amyotrophic lateral sclerosis (ALS), MAPK signaling pathway |
| MW: | 74 kDa |
| Gene Summary: | <p>This gene encodes a multifunctional protein that resides in multiple locations in the nucleus and in the cytoplasm. It interacts with a wide variety of proteins, such as apoptosis antigen Fas, centromere protein C, and transcription factor erythroblastosis virus E26 oncogene homolog 1. In the nucleus, the encoded protein functions as a potent transcription repressor that binds to sumoylated transcription factors. Its repression can be relieved by the sequestration of this protein into promyelocytic leukemia nuclear bodies or nucleoli. This protein also associates with centromeres in G2 phase. In the cytoplasm, the encoded protein may function to regulate apoptosis. The subcellular localization and function of this protein are modulated by post-translational modifications, including sumoylation, phosphorylation and polyubiquitination. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2008]</p> |

Product images:



Circular map for RC234672