

Product datasheet for **RC234842**

Bile salt activated lipase (CEL) (NM_001807) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Bile salt activated lipase (CEL) (NM_001807) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Bile salt activated lipase
Synonyms:	BAL; BSDL; BSSL; CEase; CELL; FAP; FAPP; LIPA; MODY8
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>RC234842 representing NM_001807
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGCTCACCATGGGGCGCCTGCAACTGGTTGTGTTGGCCCTCACCTGCTGCTGGGCAGTGGCGAGTGCCG
 CGAAGCTGGGCGCGTGTACACAGAAGGTGGGTTTCGTGGAAGGCGTCAATAAGAAGCTCGGCCTCCTGGG
 TGA CTCTGTGGACATCTTCAAGGCATCCCCTTCGCAGCTCCACCAAGGCCCTGGAAAACTCTCAGCCA
 CATCCTGGCTGGCAAGGGACCCTGAAGGCCAAGAATTCAAGAAGAGATGCCTGCAGGCCACCATCACCC
 AGGACAGCACCTACGGGGATGAAGACTGCCTGTACCTCAACATTTGGGTGCCCCAGGGCAGGAAGCAAGT
 CTCCCGGACCTGCCGTTATGATCTGGATCTATGGAGGCGCCTTCTCATGGGGTCCGGCCATGGGGCC
 AACTTCTCAACAACCTGTATGACGGCGAGGAGATCGCCACACGCGGAAACGTCATCGTGGTCACTT
 TCAACTACCGTGTGGCCCCCTTGGGTTCTCAGCACTGGGGACGCCAATCTGCCAGTAACTATGGCCT
 TCGGGATCAGCACATGGCCATTGCTTGGGTGAAGAGGAATATCGCGGCCCTTCGGGGGGACCCCAACAAC
 ATCACGCTCTTCGGGGAGTCTGCTGGAGGTGCCAGCGTCTCTCTGCAGACCCTCTCCCCACAACAAGG
 GCCTCATCCGGCGAGCCATCAGCCAGAGCGGCGTGGCCCTGAGTCCCTGGGTATCCAGAAAAACCCACT
 CTTCTGGGCCAAAAAGGTGGCTGAGAAGGTGGGTTGCCCTGTGGGTGATGCCGCCAGGATGGCCAGTGT
 CTGAAGGTTACTGATCCCCGAGCCCTGACGCTGGCCTATAAGGTGCCGCTGGCAGGCCTGGAGTACCCCA
 TGCTGCACTATGTGGGCTTCGTCCCTGTATTGATGGAGACTTCATCCCCGCTGACCCGATCAACCTGTA
 CGCAACGCCCGCCGACATCGACTATATAGCAGGCCAACAAACATGGACGGCCACATCTTCGCCAGCATC
 GACATGCCCTGCCATCAACAAGGGCAACAAGAAAGTCACGGAGGAGGACTTCTACAAGCTGGTCAGTGAGT
 TCACAATCACAAGGGGCTCAGAGGCGCCAAGACGACCTTTGATGTCTACACCGAGTCTGGGCCAGGA
 CCCATCCAGGAGAATAAGAAGAAGACTGTGGTGGACTTTGAGACCGATGTCTCTTCTGGTGCCACC
 GAGATTGCCCTAGCCAGCACAGAGCCAATGCCAAGAGTGCCAAGACCTACGCCTACCTGTTTTCCATC
 CCTCTCGGATGCCGCTTACCCAAATGGGTGGGGCCGACCATGCAGATGACATTAGTACGTTTTTCGG
 GAAGCCCTTCGCCACCCACAGGGCTACCGGCCCAAGACAGGACAGTCTCTAAGGCCATGATCGCCTAC
 TGGACCAACTTTGCCAAAACAGGGGACCCCAACATGGGCGACTCGGCTGTGCCACACACTGGGAACCT
 AACTACGGAAAACAGCGGCTACCTGGAGATCACCAAGAAGATGGGCAGCAGCTCCATGAAGCGGAGCCT
 GAGAACCAACTTCTGCGCTACTGGACCCTCACCTATCTGGCGCTGCCACAGTGACCGACCAGGAGGCC
 ACCCTGTGCCCCACAGGGGACTCCGAGGCCACTCCCGTGCCCCACGGGTGACTCCGAGACCGCCC
 CCGTGCCGCCACGGGTGACTCCGGGGCCCCCGTGGCGCCACGGGTGACTCCGGGGCCCCCCCCGT
 GCCGCCACGGGTGACTCCGGGGCCCCCCCCGTGCCGCCACGGGTGACTCCGGGGCCCCCCCCGTGCC
 CCCACGGGTGACTCCGGGGCCCCCCCCGTGCCGCCACGGGTGACTCCGGGGCCCCCCCCGTGCCGCCA
 CGGGTACTCCGGGGCCCCCCCCGTGCCGCCACGGGTGACTCCGGCGCCCCCCCCGTGCCGCCACGGG
 TGACGCCGGGGCCCCCCCCGTGCCGCCACGGGTGACTCCGGCGCCCCCCCCGTGCCGCCACGGGTGAC
 TCCGGGGCCCCCCCCGTGACCCACGGGTGACTCCGAGACCGCCCCGTGCCGCCACGGGTGACTCCG
 GGGCCCCCTGTGCCCCACGGGTGACTCTGAGGCTGCCCTGTGCCCCACAGATGACTCCAAGGA
 AGCTCAGATGCCTGCAGTCATTAGGTTT

ACGCGTACGCGGCGCCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC234842 representing NM_001807
 Red=Cloning site Green=Tags(s)

```
MLTMGRLQLVVLGLTCCWAVASAAKLGAVYTEGGFVEGVNKKLGLLGDSVDIFKGIPFAAPTKALENPQP
HPGWQGTLKAKNFKKRCLQATITQDSTYGDEDCLYLNIVWPQGRKQVSRDLPVMIWIYGGAFMLGSGHGA
NFLNNLYLDGEEIATRGNVIVVTFNYRVGPLGFLSTGDANLPNGYGLRDQHMAIAWVKRNIAAFGGDPNN
ITLFGESAGGASVSLQTLSPYKGLIRRAISQSGVALSPWVIQKNPLFWAKKVAEKVGCPCVGDAAARMAQC
LKVTDPRALTLAYKVPLAGLEYPMLHYVGFVVIDGDFIPADPINLYANAADIDYIAGTNNMDGHIFASI
DMPAINKGNKKVTEEDFYKLVSEFTITKGLRGAKTTFDVYTESWAQDPSQENKKTVVDFETDVLFLVPT
EIALAQHRANAKSAKTYAYLFSHPSRMPVYPKWVGADHADDIQYVFGKPFATPTGYRQDRTVSKAMIAIY
WTNFAKTGDPNMGDSAVPTHWEPTYTENSYLEITKKMGSSSMKRSLRTNFLRYWTLTYLALPTVTDQEA
TPVPPTGDSEATVPVPTGDSEATVPVPTGDSGAPPVPTGDSGAPPVPTGDSGAPPVPTGDSGAPPVPT
PTGDSGAPPVPTGDSGAPPVPTGDSGAPPVPTGDSGAPPVPTGDSGAPPVPTGDAGPPPVPPTGDSGAPPVPTGD
SGAPPVPTGDSEATVPVPTGDSGAPPVPTGDSEAAPVPTDDSKAQMPAVIRF
```



TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_001807

ORF Size: 2268 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001807.4](#), [NP_001798.2](#)

RefSeq Size: 2386 bp

RefSeq ORF: 2262 bp

Locus ID: 1056

Cytogenetics: 9q34.13


Protein Families: Druggable Genome

Protein Pathways: Glycerolipid metabolism, Metabolic pathways, Steroid biosynthesis

MW: 80.1 kDa

Gene Summary: The protein encoded by this gene is a glycoprotein secreted from the pancreas into the digestive tract and from the lactating mammary gland into human milk. The physiological role of this protein is in cholesterol and lipid-soluble vitamin ester hydrolysis and absorption. This encoded protein promotes large chylomicron production in the intestine. Also its presence in plasma suggests its interactions with cholesterol and oxidized lipoproteins to modulate the progression of atherosclerosis. In pancreatic tumoral cells, this encoded protein is thought to be sequestered within the Golgi compartment and is probably not secreted. This gene contains a variable number of tandem repeat (VNTR) polymorphism in the coding region that may influence the function of the encoded protein. [provided by RefSeq, Jul 2008]

Product images:



Circular map for RC234842