

Product datasheet for **RC235518**

Proteasome subunit beta type 2 (PSMB2) (NM_001199780) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Proteasome subunit beta type 2 (PSMB2) (NM_001199780) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: PSMB2
Synonyms: HC7-I
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
ORF Nucleotide Sequence: >RC235518 representing NM_001199780
Red=Cloning site Blue=ORF Green=Tags(s)

TTTGTAAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**GCGATCGCC**

ATGGACTACCTGGCAGCCTTGGCCAAGGCCCTTTGCAGCCACGGCTATGGTGCCTTCTGACTCTCA
GTATCCTCGACCGATACTACACACCGACTATCTCACGTGAGAGGGCAGTGAACCTTAGGAAATGTCT
GGAGGAGCTCCAGAAACGCTTCATCCTGAATCTGCCAACCTTCAGTGTTCAATCATTGACAAAAATGGC
ATCCATGACCTGGATAACATTTCTTCCCAAACAGGGCTCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC235518 representing NM_001199780
Red=Cloning site Green=Tags(s)

MDYLAALAKAPFAAHGYGAFLTLSILDYYTPTISRERAVELLRKCLEELQKRFILNLPTFSVRIIDKNG
IHDLDNISFPKQGS


TRTRPLEQK**L**ISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-MluI



[View online »](#)

Cloning Scheme:

Plasmid Map:


ACCN: NM_001199780

ORF Size: 252 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

| | |
|-------------------------------|--|
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | <ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | <u>NM_001199780.1, NP_001186709.1</u> |
| RefSeq Size: | 4235 bp |
| RefSeq ORF: | 255 bp |
| Locus ID: | 5690 |
| Cytogenetics: | 1p34.3 |
| Protein Families: | Druggable Genome, Protease |
| Protein Pathways: | Proteasome |
| MW: | 10 kDa |
| Gene Summary: | <p>The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Dec 2010]</p> |