

Product datasheet for RC235521

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Apc11 (ANAPC11) (NM_001289416) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Apc11 (ANAPC11) (NM 001289416) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: Apc11

Synonyms: APC11; Apc11p; HSPC214

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >RC235521 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

AGCAGCACTGCCCCATGTGCCGCCAGGAATGGAAGTTCAAGGAG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC235521 protein sequence

Red=Cloning site Green=Tags(s)

X*R*RLSAGTAWPLGSGWPTMRTVASAGWHLTDAALTARCPATTARWCGASAPTASTCIASSSGCTHSRC

SSTAPCAARNGSSR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6486 c01.zip


Restriction Sites: Sgfl-Mlul





Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM 001289416

ORF Size: 252 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001289416.1</u>, <u>NP 001276345.1</u>

RefSeq Size: 914 bp RefSeq ORF: 255 bp Locus ID: 51529



UniProt ID: Q9NYG5

Cytogenetics: 17q25.3

Protein Families: Druggable Genome

Protein Pathways: Cell cycle, Oocyte meiosis, Progesterone-mediated oocyte maturation, Ubiquitin mediated

proteolysis


MW: 9.8 kDa

Gene Summary: Together with the cullin protein ANAPC2, constitutes the catalytic component of the anaphase

promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains. May recruit the E2 ubiquitin-conjugating

enzymes to the complex.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RC235521