

## Product datasheet for **RC235656**

### **CXCL11 (NM\_001302123) Human Tagged ORF Clone**

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** CXCL11 (NM\_001302123) Human Tagged ORF Clone  
**Tag:** Myc-DDK  
**Symbol:** CXCL11  
**Synonyms:** b-R1; H174; I-TAC; IP-9; IP9; SCYB9B; SCYB11  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Cell Selection:** Neomycin  
**ORF Nucleotide Sequence:** >RC235656 representing NM\_001302123  
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**GCGATCGCC**

ATGAGTGTGAAGGGCATGGCTATAGCCTTGGCTGTGATATTGTGTGCTACAGTTGTTCAAGGCTTCCCCA  
TGTTCAAAGAGGACGCTGTCTTTGCATAGGCCCTGGGGTAAAAGCAGTAAAAGTGGCAGATATTGAGAA  
AGCCTCCATAATGTACCAAGTAACAACGTGACAAAATAGAAGTGATTATTACCCTGAAAGAAAATAAA  
GGACAACGATGCCTAAATCCCAAATCGAAGCAAGCAAGGCTTATAATCAAATTGAAAGAAAAGAAATTTTAA  
AAAATATCAAACATATGAAGTCCTGGAAAAGAGCATC

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >RC235656 representing NM\_001302123  
Red=Cloning site Green=Tags(s)

MSVKGMAIALAVILCATVVQGFPMFKRGRCLCIGPGVKAVKVADIEKASIMYPSNNCDKIEVIITLKENK  
GQRCLNPKSKQARLIIKLERIFKNIKTYEVLEKSI


**TR**TRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** Sgfl-MluI



[View online »](#)

**Cloning Scheme:**

**Plasmid Map:**


ACCN: NM\_001302123

ORF Size: 318 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001302123.2</a>
<b>RefSeq Size:</b>	1603 bp
<b>RefSeq ORF:</b>	321 bp
<b>Locus ID:</b>	6373
<b>UniProt ID:</b>	<a href="#">O14625</a>
<b>Cytogenetics:</b>	4q21.1
<b>Protein Families:</b>	Druggable Genome, Secreted Protein, Transmembrane
<b>Protein Pathways:</b>	Chemokine signaling pathway, Cytokine-cytokine receptor interaction, Toll-like receptor signaling pathway
<b>MW:</b>	12.2 kDa
<b>Gene Summary:</b>	Chemokines are a group of small (approximately 8 to 14 kD), mostly basic, structurally related molecules that regulate cell trafficking of various types of leukocytes through interactions with a subset of 7-transmembrane, G protein-coupled receptors. Chemokines also play fundamental roles in the development, homeostasis, and function of the immune system, and they have effects on cells of the central nervous system as well as on endothelial cells involved in angiogenesis or angiostasis. Chemokines are divided into 2 major subfamilies, CXC and CC. This antimicrobial gene is a CXC member of the chemokine superfamily. Its encoded protein induces a chemotactic response in activated T-cells and is the dominant ligand for CXC receptor-3. The gene encoding this protein contains 4 exons and at least three polyadenylation signals which might reflect cell-specific regulation of expression. IFN-gamma is a potent inducer of transcription of this gene. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2014]