

Product datasheet for RC235659

MARK1 (NM 001286129) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: MARK1 (NM_001286129) Human Tagged ORF Clone

Tag: Myc-DDK
Symbol: MARK1

Synonyms: MARK; Par-1c; Par1c

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

ORF Nucleotide >RC235659 representing NM_001286129
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

CTCAGCTAAATCCTACCAGTCTACAAAAGGTATTTAAT

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC235659 representing NM_001286129

Red=Cloning site Green=Tags(s)

MSARTPLPTVNERDTENHTSVDGYTEPHIQPTKSSSRQNIPRCRNSITSATDEQPHIGNYRLQKTIGKGN

FAKVKLARHVLTGREVAVKIIDKTQLNPTSLQKVFN

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-Mlul



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

Plasmid Map:



ACCN: NM_001286129

ORF Size: 318 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info



MARK1 (NM_001286129) Human Tagged ORF Clone - RC235659

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001286129.2</u>

RefSeq Size: 2747 bp
RefSeq ORF: 321 bp
Locus ID: 4139
Cytogenetics: 1q41

Protein Families: Druggable Genome, Protein Kinase

MW: 12.3 kDa

Gene Summary: Serine/threonine-protein kinase (PubMed:23666762). Involved in cell polarity and microtubule

dynamics regulation. Phosphorylates DCX, MAP2 and MAP4. Phosphorylates the microtubule-

associated protein MAPT/TAU (PubMed:23666762). Involved in cell polarity by

phosphorylating the microtubule-associated proteins MAP2, MAP4 and MAPT/TAU at KXGS motifs, causing detachment from microtubules, and their disassembly. Involved in the regulation of neuronal migration through its dual activities in regulating cellular polarity and microtubule dynamics, possibly by phosphorylating and regulating DCX. Also acts as a

positive regulator of the Wnt signaling pathway, probably by mediating phosphorylation of

dishevelled proteins (DVL1, DVL2 and/or DVL3).[UniProtKB/Swiss-Prot Function]