

Product datasheet for RC235675

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Fucose mutarotase (FUOM) (NM_001301828) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Fucose mutarotase (FUOM) (NM_001301828) Human Tagged ORF Clone

Tag: Myc-DDK
Symbol: FUOM

Synonyms:C10orf125; FucM; FUCUVector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

ORF Nucleotide >RC235675 representing NM_001301828
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

GAAACCTCATCCTCAGGAAGGGGGTGCTTGCCCTCAACCCCCTGCTG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC235675 representing NM_001301828

Red=Cloning site Green=Tags(s)

MEIRADGLGIPQLLEAVLKLLPLDTYVESPAAVMELVPSDKERGLQTPVWTEYESILRRAGCVRALAKIE

RFEFYERAKKAFAVVATGETALYGNLILRKGVLALNPLL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Restriction Sites: Sgfl-Mlul






Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

Plasmid Map:



ACCN: NM_001301828

ORF Size: 327 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info



Fucose mutarotase (FUOM) (NM_001301828) Human Tagged ORF Clone - RC235675

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001301828.2</u>

 RefSeq Size:
 906 bp

 RefSeq ORF:
 330 bp

 Locus ID:
 282969

 UniProt ID:
 A2VDF0

 Cytogenetics:
 10q26.3

 MW:
 12.6 kDa

Gene Summary: Involved in the interconversion between alpha- and beta-L-fucoses. L-Fucose (6-deoxy-L-

galactose) exists as alpha-L-fucose (29.5%) and beta-L-fucose (70.5%), the beta-form is metabolized through the salvage pathway. GDP-L-fucose formed either by the de novo or salvage pathways is transported into the endoplasmic reticulum, where it serves as a substrate for N- and O-glycosylations by fucosyltransferases. Fucosylated structures

expressed on cell surfaces or secreted in biological fluids are believed to play a critical role in

cell-cell adhesion and recognition processes.[UniProtKB/Swiss-Prot Function]