

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for RC235890

### CRSP8 (MED27) (NM\_001253882) Human Tagged ORF Clone

## **Product data:**

Product Type:	Expression Plasmids
Product Name:	CRSP8 (MED27) (NM_001253882) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	CRSP8
Synonyms:	CRAP34; CRSP8; CRSP34; MED3; NEDSCAC; TRAP37
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	<pre>&gt;RC235890 representing NM_001253882 Red=Cloning site Blue=ORF Green=Tags(s)</pre>
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGGCGGACGTGATAAATGTCAGTGTGAACCTGGAGGCCTTTTCCCAGGCCATTAGTGCCATCCAGGCGC TGCGCTCCAGCGTGAGCAGGGTGTTCGACTGCCTGAAGGATGGGATGCGGAACAAGGAGACGCTGGAGGG CCGGGAGAAGGCCTTTATTGCGCACTTCCAGGACAACTTACATTCGGTCAACCGGGACCTCAATGAGCTG GAACGTCTGAGCAATCTGGTAGGCAAGCCATCTGAGAACCATCCTCTTCATAACAGTGGGCTGTTAAGCC TGGATCCTGTGCAGGACAAAACTCCTCTCTATAGTCAACTCCTTCAAGCATATAAGTGGTCAAACAAGAA AGAGCAACTTTCTATTCCAAGAATATTCCATTGGAAAGTC
	ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAG <b>GTTTAA</b>
Protein Sequence:	>RC235890 representing NM_001253882 <mark>Red</mark> =Cloning site Green=Tags(s)
	MADVINVSVNLEAFSQAISAIQALRSSVSRVFDCLKDGMRNKETLEGREKAFIAHFQDNLHSVNRDLNEL ERLSNLVGKPSENHPLHNSGLLSLDPVQDKTPLYSQLLQAYKWSNKKEQLSIPRIFHWKV
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
<b>Restriction Sites:</b>	Sgfl-Mlul



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

#### **Cloning Scheme:**



\* The last codon before the Stop codon of the ORF


ACCN:	NM_001253882
ORF Size:	390 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM 001253882.2</u>
RefSeq Size:	799 bp

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	CRSP8 (MED27) (NM_001253882) Human Tagged ORF Clone – RC235890	
RefSeq ORF:	393 bp	
Locus ID:	9442	
UniProt ID:	<u>Q6P2C8</u>	
Cytogenetics:	9q34.13	
MW:	15.3 kDa	

Gene Summary: The activation of gene transcription is a multistep process that is triggered by factors that recognize transcriptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 5. [provided by RefSeq, Dec 2011]

# **Product images:**



Circular map for RC235890

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US