

## Product datasheet for **RC236158**

### GDNF (NM\_001278098) Human Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** GDNF (NM\_001278098) Human Tagged ORF Clone  
**Tag:** Myc-DDK  
**Symbol:** GDNF  
**Synonyms:** ATF; ATF1; ATF2; HFB1-GDNF; HSCR3  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Cell Selection:** Neomycin  
**ORF Nucleotide Sequence:** >RC236158 representing NM\_001278098  
**Red=Cloning site Blue=ORF Green=Tags(s)**

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**GCGATCGCC**

ATGCCAGAGGATTATCCTGATCAGTTCGATGATGTCATGGATTTTATTCAAGCCACCATTAAAAGACTGA  
AAAGGTCACCAGATAAACAAATGGCAGTGCTTCTAGAAGAGAGCGGAATCGGCAGGCTGCAGCTGCCAA  
CCCAGAGAATCCAGAGGAAAAGGTCGGAGAGGCCAGAGGGGCAAAAACCGGGTTGTGTCTTAAGTCA  
ATACATTTAAATGTCAGTCACTGACTTGGGTCTGGGCTATGAAACCAAGGAGGAAGTATTTTAGTACTGCA  
GCGGCTCTTGCATGCAGCTGAGACAACGTACGACAAAATATTGAAAACTTATCCAGAAATAGAAGGCT  
GGTGAGTGACAAAGTAGGGCAGGCATGTTGCAGACCCATCGCCTTTGATGATGACCTGCTGTTTTAGAT  
GATAACCTGGTTTACCATATTCTAAGAAAGCATTCCGCTAAAAGGTGGATGTATC

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >RC236158 representing NM\_001278098  
**Red=Cloning site Green=Tags(s)**

MPEDYPDQFDDVMDFIQATIKRLKRSPDKQMAVLP RRERNRQAAAAANPENS RKGRRGRGKNRGCVLTA  
IHLNVTDLGLGYETKEELIFRYCSGSCDAAETTYDKILKNLSRNRRLVSDKVGQACCRPIAFDDDL SFLD  
DNLVYHILRKHSKRKCGCI

**TRTRPLEQKLI**SEEDLAANDILDYKDDDDKV

**Restriction Sites:** Sgfl-Mlul




[View online »](#)

**Cloning Scheme:**



**Plasmid Map:**



**ACCN:** NM\_001278098

**ORF Size:** 477 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u>NM_001278098.1, NP_001265027.1</u>
<b>RefSeq Size:</b>	3638 bp
<b>RefSeq ORF:</b>	480 bp
<b>Locus ID:</b>	2668
<b>UniProt ID:</b>	<u>P39905</u>
<b>Cytogenetics:</b>	5p13.2
<b>Protein Families:</b>	Druggable Genome, Secreted Protein, Transmembrane
<b>MW:</b>	18.6 kDa
<b>Gene Summary:</b>	<p>This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. The recombinant form of this protein, a highly conserved neurotrophic factor, was shown to promote the survival and differentiation of dopaminergic neurons in culture, and was able to prevent apoptosis of motor neurons induced by axotomy. This protein is a ligand for the product of the RET (rearranged during transfection) protooncogene. Mutations in this gene may be associated with Hirschsprung disease and Tourette syndrome. This gene encodes multiple protein isoforms that may undergo similar proteolytic processing. [provided by RefSeq, Aug 2016]</p>