

Product datasheet for RC236351

FPGT (NM_001199329) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	FPGT (NM_001199329) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	FPGT
Synonyms:	GFPP
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RC236351 representing NM_001199329 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGCGTGCTGTGCGGCGCGGTCTCAGGGAAGGTGGGGCTATGGCAGCTGCTAGGGACCTCCGGAAGTAT
 CGCTGCGAGAAGCCACCCAGCGAAAATTGCGGAGGTTTTCCGAGCTAAGAGGCAAACCTGTAGCACGTGG
 AGAATTCTGGGACATAGTTGCAATAACAGCGGCTGATGAAAAACAGGAACCTGCTTACAACCAACAGCTG
 TCAGAAAAGCTGAAAAGAAAGGAGTTACCCCTTGGAGTTCAATATCACGTTTTTGTGGATCCTGCTGGAG
 CCAAAATTGGAATGGAGGATCAACACTTTGTGCCCTTCAATGTTTGGAAAAGCTATATGGAGATAAATG
 GAATTCCTTTTACCATCTTATTAATTCACCTCTGTATCTTTCCAGATATACCAGAATGCTCTGGCAAAACAT
 CCTGTATCATTCAAAGCATACTGGATTCAAGATGTTCTGTGGCACCTGGCTCAGTTGTGGAGTATTCCAG
 ATTGGGGCCTGATGTTTCAGTTGGGAAAACCTGCATTATTAGTGGTTCTTACATCC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:	>RC236351 representing NM_001199329 Red=Cloning site Green=Tags(s)
-------------------	---

MRAVRRGLREGGAMAAARDPPEVSLREATQRKLRRFSELRGKLVARGEFDIVAITAADEKQELAYNQQL
 SEKLKRKELPLGVQYHVFDVPAGAKIGNGGSTLCALQCLEKLYGDKWNSFTILLIHSVSFQIYQNALAKH
 PVSFKAYWIQDVLWHLAQLWSIPDWGLMFQLGKTALLVVLTS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:	Sgfl-MluI
--------------------	-----------


[View online »](#)

Cloning Scheme:


ACCN: NM_001199329

ORF Size: 546 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

RefSeq: [NM_001199329.2](#), [NP_001186258.2](#)


RefSeq Size: 3905 bp

RefSeq ORF: 510 bp

Locus ID: 8790

UniProt ID:	<u>O14772</u>
Cytogenetics:	1p31.1
Protein Pathways:	Amino sugar and nucleotide sugar metabolism, Fructose and mannose metabolism, Metabolic pathways
MW:	21 kDa
Gene Summary:	L-fucose is a key sugar in glycoproteins and other complex carbohydrates since it may be involved in many of the functional roles of these macromolecules, such as in cell-cell recognition. The fucosyl donor for these fucosylated oligosaccharides is GDP-beta-L-fucose. There are two alternate pathways for the biosynthesis of GDP-fucose; the major pathway converts GDP-alpha-D-mannose to GDP-beta-L-fucose. The protein encoded by this gene participates in an alternate pathway that is present in certain mammalian tissues, such as liver and kidney, and appears to function as a salvage pathway to reutilize L-fucose arising from the turnover of glycoproteins and glycolipids. This pathway involves the phosphorylation of L-fucose to form beta-L-fucose-1-phosphate, and then condensation of the beta-L-fucose-1-phosphate with GTP by fucose-1-phosphate guanylyltransferase to form GDP-beta-L-fucose. Alternative splicing results in multiple transcript variants. Read-through transcription also exists between this gene and the neighboring downstream TNNI3 interacting kinase (TNNI3K) gene. [provided by RefSeq, Dec 2010]

Product images:



Circular map for RC236351