

# Product datasheet for RC236473

## MICA (NM 001289154) Human Tagged ORF Clone

### **Product data:**

**Product Type: Expression Plasmids** 

**Product Name:** MICA (NM\_001289154) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: MICA

Synonyms: MIC-A; PERB11.1

Vector: pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

Cell Selection: Neomycin

**ORF Nucleotide** >RC236473 representing NM\_001289154 Red=Cloning site Blue=ORF Green=Tags(s) Sequence:

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

**GCCGCGATCGCC** 

ATGGGACAGAGAGACCAGGGACTTGACAGGGAACGGAAAGGACCTCAGGATGACCCTGGCTCATATCAAG AGACTGCCTGCAGGAACTACGGCGATATCTAGAATCCGGCGTAGTCCTGAGGAGAACAGTGCCCCCCATG GTGAATGTCACCCGCAGCGAGGCCTCAGAGGGCAACATCACCGTGACATGCAGGGCTTCCAGCTTCTATC CCCGGAATATCATACTGACCTGGCGTCAGGATGGGGTATCTTTGAGCCACGACACCCAGCAGTGGGGGGA TGTCCTGCCTGATGGGAATGGAACCTACCAGACCTGGGTGGCCACCAGGATTTGCCGAGGAGAGGAGCAG AGGTTCACCTGCTACATGGAACACAGCGGGAATCACAGCACTCACCCTGTGCCCTCTGGGAAAGTGCTGG TGCTTCAGAGTCATTGGCAGACATTCCATGTTTCTGCTGTTGCTGCTGCTGCTGCTATTTTTGTTATTA

TTATTTTCTATGTCCGTTGTTG

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >RC236473 representing NM\_001289154

Red=Cloning site Green=Tags(s)

MGQRDQGLDRERKGPQDDPGSYQGPERRNFLKEDAMKTKTHYHAMHADCLQELRRYLESGVVLRRTVPPM VNVTRSEASEGNITVTCRASSFYPRNIILTWRQDGVSLSHDTQQWGDVLPDGNGTYQTWVATRICRGEEQ

RFTCYMEHSGNHSTHPVPSGKVLVLQSHWQTFHVSAVAAGCCYFCYYYFLCPLL

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

**Restriction Sites:** Sgfl-Mlul



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

### Plasmid Map:



**ACCN:** NM\_001289154

ORF Size: 582 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** <u>NM 001289154.2</u>

 RefSeq Size:
 2086 bp

 RefSeq ORF:
 585 bp

 Locus ID:
 100507436

 Cytogenetics:
 6p21.33

 MW:
 22.7 kDa

Gene Summary: This gene encodes the highly polymorphic major histocompatability complex class I chain-

related protein A. The protein product is expressed on the cell surface, although unlike canonical class I molecules it does not seem to associate with beta-2-microglobulin. It is a ligand for the NKG2-D type II integral membrane protein receptor. The protein functions as a stress-induced antigen that is broadly recognized by intestinal epithelial gamma delta T cells. Variations in this gene have been associated with susceptibility to psoriasis 1 and psoriatic arthritis, and the shedding of MICA-related antibodies and ligands is involved in the progression from monoclonal gammopathy of undetermined significance to multiple

myeloma. Alternative splicing of this gene results in multiple transcript variants. [provided by

RefSeq, Jan 2014]