

Product datasheet for RC236660

RFC2 (NM 001278793) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: RFC2 (NM_001278793) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: RFC2

Synonyms: RFC40

Vector: pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

Cell Selection: Neomycin

ORF Nucleotide >RC236660 representing NM_001278793
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGTTGGAACTCAATGCTTCAAATGACAGCATGACCGACGAGCCCAGCAAGCCTTGAGGAGAACCATGG
AAATCTACTCTAAAACCACTCGCTTCGCCCTTGCTTGTAATGCTTCGGATAAGATCATCGAGCCCATTCA
GTCCCGCTGTGCAGTCCTCCGGTACACAAAGCTGACCGACGCCCAGATCCTCACCAGGCTGATGAATGTT
ATCGAGAAGGAGAGGGTACCCTACACTGATGACGGCCTAGAAGCCATCATCTTCACGGCCCAGGGAGACA
TGAGGCCAGGCGCTGAACAACCTGCAGTCCACCTTCTCAGGATTTGGCTTCATTAACAGTGAGAACGTGTT
CAAGGTCTGTGACGAGCCCCACCCACTGCTGGTAAAGGAGATGATCCAGCACTGTGTGAATGCCAACATT
GACGAAGCCTACAAGATTCTTGCTCACTTGTGGCATCTGGGCTACTCACCAGAAGATATCATTGGCAACA
TCTTTCGAGTGTGTAAAACTTTCCAAATGGCAGAATACCTGAAACTGGAGTTTATCAAGGAAATTGGATA
CACTCACATGAAAAATAGCGGAAGGAGTGAACTCTCTTTTTGCAGATGGCAGGCCTCCTGGCAAGGCTGTGT
CAGAAGACAATGGCCCCGGTGGCCAGT

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >RC236660 representing NM_001278793

Red=Cloning site Green=Tags(s)


MLELNASNDSMTDGAQQALRRTMEIYSKTTRFALACNASDKIIEPIQSRCAVLRYTKLTDAQILTRLMNV IEKERVPYTDDGLEAIIFTAQGDMRQALNNLQSTFSGFGFINSENVFKVCDEPHPLLVKEMIQHCVNANI DEAYKILAHLWHLGYSPEDIIGNIFRVCKTFQMAEYLKLEFIKEIGYTHMKIAEGVNSLLQMAGLLARLC QKTMAPVAS


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

Sgfl-Mlul


Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

Plasmid Map:



ACCN: NM_001278793

ORF Size: 657 bp

RFC2 (NM_001278793) Human Tagged ORF Clone - RC236660

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001278793.1, NP 001265722.1</u>

 RefSeq Size:
 1759 bp

 RefSeq ORF:
 660 bp

 Locus ID:
 5982

 UniProt ID:
 P35250

Cytogenetics: 7q11.23

Protein Families: Druggable Genome, Stem cell - Pluripotency

Protein Pathways: DNA replication, Mismatch repair, Nucleotide excision repair

MW: 25.2 kDa

Gene Summary: This gene encodes a member of the activator 1 small subunits family. The elongation of

primed DNA templates by DNA polymerase delta and epsilon requires the action of the accessory proteins, proliferating cell nuclear antigen (PCNA) and replication factor C (RFC). Replication factor C, also called activator 1, is a protein complex consisting of five distinct subunits. This gene encodes the 40 kD subunit, which has been shown to be responsible for binding ATP and may help promote cell survival. Disruption of this gene is associated with Williams syndrome. Alternatively spliced transcript variants encoding distinct isoforms have been described. A pseudogene of this gene has been defined on chromosome 2. [provided by

RefSeq, Jul 2013]