

Product datasheet for RC236663

WFDC1 (NM 001282467) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: WFDC1 (NM_001282467) Human Tagged ORF Clone

Tag: Myc-DDK
Symbol: WFDC1
Synonyms: PS20

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

ORF Nucleotide >RC236663 representing NM_001282467
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAGGTTTAA



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >RC236663 representing NM_001282467

Red=Cloning site Green=Tags(s)


MPLTGVGPGSCRRQIIRALCLLLLLLHAGSAKNIWKRALPARLAEKSRAEEAGAPGGPRQPRADRCPPPP RTLPPGACQAARCQADSECPRHRRCCYNGCAYACLEAVPPPPVLDWLVQPKPRWLGGNGWLLDGPEEVLQ EACSTTEDGAEPLLCPSGYECHILSPGDVAEGIPNRGQCVKQRRQADGRILRHKLYKEYPEGDSKNVAEP GRGQQKHFQ


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

Sgfl-Mlul


Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

Plasmid Map:



ACCN: NM_001282467

ORF Size: 657 bp



WFDC1 (NM_001282467) Human Tagged ORF Clone - RC236663

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

16q24.1

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001282467.2</u>

 RefSeq Size:
 1472 bp

 RefSeq ORF:
 660 bp

 Locus ID:
 58189

 UniProt ID:
 Q9HC57

Protein Families: Secreted Protein

MW: 24.4 kDa

Cytogenetics:

Gene Summary: This gene encodes a member of the WAP-type four disulfide core domain family. The WAP-

type four-disulfide core domain contains eight cysteines forming four disulfide bonds at the core of the protein, and functions as a protease inhibitor in many family members. This gene is mapped to chromosome 16q24, an area of frequent loss of heterozygosity in cancers, including prostate, breast and hepatocellular cancers and Wilms' tumor. This gene is downregulated in many cancer types and may be involved in the inhibition of cell

proliferation. The encoded protein may also play a role in the susceptibility of certain CD4 memory T cells to human immunodeficiency virus infection. Alternative splicing results in

multiple transcript variants. [provided by RefSeq, Sep 2013]