

Product datasheet for **RC236672**

PAX5 (NM_001280551) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: PAX5 (NM_001280551) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: PAX5
Synonyms: ALL3; BSAP
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
ORF Nucleotide Sequence: >RC236672 representing NM_001280551
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**GCGATCGCC**

ATGTTTGCCTGGGAGATCAGGGACCGGCTGCTGGCAGAGCGGGTGTGTGACAATGACACCGTGCCTAGCG
TCAGTTCCATCAACAGGATCATCCGGACAAAAGTACAGCAGCCACCAACCAACCAGTCCCAGTCCAG
TCACAGCATAGTGTCCACTGGCTCCGTGACGCAGGTGTCTCGGTGAGCAGGATTCCGGCCGGCTCGTCG
TACTCCATCAGCGGCATCCTGGGCATCAGTCCCCAGCGCCGACACCAACAAGCGCAAGAGAGACGGAAG
GTATTCAGGAGTCTCCGGTGCCGAACGGCCACTCGCTTCCGGGCAGAGACTTCTCCGGAAGCAGATGCG
GGGAGACTTGTTACACAGCAGCAGCTGGAGGTGCTGGACCGCGTGTGAGAGGCAGCACTACTCAGAC
ATCTTCACCACCACAGAGCCCATCAAGCCCAGCAGACCACAGAGTATTCAGCCATGGCCTCGCTGGCTG
GTGGGCTGGACGACATGAAGGCCAATCTGGCCAGCCCCACCCCTGCTGACATCGGGAGCAGTGTGCCAGG
CCCGCAGTCTACCCATTGTGACAGGCTCCCCCTACTATTATAGCGCTGCCGCCGAGGAGCCGCCCA
CCTGCAGCCGCCACTGCCTATGACCGTCAC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >RC236672 representing NM_001280551
 Red=Cloning site Green=Tags(s)

MFAWEIRDRLLAERVCDNDTVPSVSSINRIIRTKVQPPNPVPPASSHSIVSTGVSVTQVSSVSTDSAGSS
 YSISGILGITSPSADTNKRKRDEGIQESVPVNGHSLPGRDFLRKQMRGDLFTQQQLEVLDRVFERQHYS
 IFTTTEPIKPEQTTEYSAMASLAGGLDDMKANLASPTPADIGSSVPGPQSYPIVGTSPYYSAARGAAP
 PAAATAYDRH


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_001280551

ORF Size: 660 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001280551.2
RefSeq Size:	8554 bp
RefSeq ORF:	663 bp
Locus ID:	5079
Cytogenetics:	9p13.2
Protein Families:	Transcription Factors
MW:	24.1 kDa
Gene Summary:	This gene encodes a member of the paired box (PAX) family of transcription factors. The central feature of this gene family is a novel, highly conserved DNA-binding motif, known as the paired box. Paired box transcription factors are important regulators in early development, and alterations in the expression of their genes are thought to contribute to neoplastic transformation. This gene encodes the B-cell lineage specific activator protein that is expressed at early, but not late stages of B-cell differentiation. Its expression has also been detected in developing CNS and testis and so the encoded protein may also play a role in neural development and spermatogenesis. This gene is located at 9p13, which is involved in t(9;14)(p13;q32) translocations recurring in small lymphocytic lymphomas of the plasmacytoid subtype, and in derived large-cell lymphomas. This translocation brings the potent E-mu enhancer of the IgH gene into close proximity of the PAX5 promoter, suggesting that the deregulation of transcription of this gene contributes to the pathogenesis of these lymphomas. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2013]